

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12Q 1/68, C12P 19/34 C07H 21/02, 21/04, C07K 13/00	A1	(11) International Publication Number: WO 94/09162 (43) International Publication Date: 28 April 1994 (28.04.94)
(21) International Application Number: PCT/US93/10072 (22) International Filing Date: 20 October 1993 (20.10.93) (30) Priority data: 07/963,723 20 October 1992 (20.10.92) US (71) Applicant: EMORY UNIVERSITY SCHOOL OF MEDICINE [US/US]; Atlanta, GA 30322 (US). (72) Inventor: WALLACE, Douglas, C. ; 2506 Echo Drive, Atlanta, GA 30345 (US). (74) Agents: NEEDLE, William, H. et al.; Needle & Rosenberg, 127 Peachtree Street, N.E., Suite 1200, Atlanta, GA 30303 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: DETECTION OF MITOCHONDRIAL DNA MUTATIONS ASSOCIATED WITH ALZHEIMER'S DISEASE AND PARKINSON'S DISEASE (57) Abstract This invention provides a method of diagnosing or predicting a predisposition to Alzheimer's disease and/or Parkinson's disease. The method comprises detecting in a sample from a subject the presence of a mutation, for example, in nucleotide position 4,336, 3,397, 3,196 or an insertion between positions 956 and 965, of mitochondrial DNA. The presence of the mutation indicates the presence of or a predisposition to Alzheimer's disease. Since each mutation increases the likelihood of developing or having Alzheimer's disease, the detection of more than one of the mutations in an individual can increase the probability of having or developing the disease. The invention also provides a method of determining mutations associated with the presence of or predisposition to Alzheimer's and/or Parkinson's disease. The method comprises: a) obtaining a mitochondrial DNA-containing sample from a subject with Alzheimer's disease; b) determining the presence of mutations in the mitochondrial DNA; c) comparing the mutations found in normal subject; and d) determining which mutations have a greater rate of occurrence in the subject with Alzheimer's disease.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CP	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TC	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

DETECTION OF MITOCHONDRIAL DNA MUTATIONS ASSOCIATED WITH
ALZHEIMER'S DISEASE AND PARKINSON'S DISEASE

BACKGROUND OF THE INVENTION

5

Alzheimer's disease is a progressive neurodegenerative disorder characterized clinically by cognitive decline with onset usually after the age of 60 years. Its major neuropathological features include senile plaques, many with abnormal neurites (neuritic plaques),
10 neurofibrillary tangles within neuronal perikarya, and amyloid angiopathy. Several neurotransmitter systems are perturbed in Alzheimer's disease, with cholinergic deficiency being most prominent and associated with cell loss in the nucleus basalis of Meynert¹.

15

Substantial clinical, pathological, biochemical, and genetic heterogeneity exists in Alzheimer's disease. Frequently, Alzheimer's disease patients exhibit extrapyramidal signs of Parkinson's disease^{2,3} and have coexisting neuropathological features of Parkinson's disease (AD+PD). These can include substantia nigra degeneration and Lewy
20 bodies in the pigmented nuclei and nucleus basalis of about 20-40% of neuropathologically confirmed autopsy brains^{4,5}. Cases displaying cortical Lewy bodies have been called "diffuse Lewy body disease"⁶⁻¹⁰, and are increasingly recognized with and without concomitant Alzheimer's disease pathology. Thus, the clinical and
25 neuropathological features of Alzheimer's disease and Parkinson's disease overlap extensively, suggesting that they may represent a spectrum of disease with related causal mechanisms.

30

Our recent investigations have suggested that oxidative phosphorylation (OXPHOS) defects may play a role in the pathogenesis of Alzheimer's disease and Parkinson's disease^{11,12}. OXPHOS enzyme assays in Parkinson's disease brains have shown Complex I defects¹³⁻¹⁵ as well as systemic OXPHOS defects in platelets¹⁶ and skeletal muscle^{17,18}. Similarly, Alzheimer's disease patients have shown Complex
35 IV defects in platelets¹⁹ and abnormalities in mitochondrial respiration in neocortex²⁰ and fibroblasts²¹. Treatment of normal human fibroblasts with the OXPHOS uncoupler CCCP (carbonyl cyanide m-chlorophenylhydrazine) results in a 10-fold increase in the proportion of cells reacting with an antibody to paired helical filaments and a

157-fold increase in cells reacting to the Alzheimer's monoclonal antibody-50²², suggesting a linkage between OXPHOS defects and Alzheimer's disease pathology. Abnormal mitochondria with paracrystalline inclusions, like those frequently encountered in the muscle from patients with mitochondrial DNA deletions and point mutations²³, have been described in the brain of a patient with AD+PD pathology²⁴ and patients with Alzheimer's disease²⁵.

OXPHOS is composed of five enzyme complexes assembled from 13 mitochondrial DNA (mtDNA) and approximately 50 nuclear DNA subunits. Complex I (NADH:ubiquinone oxidoreductase) contains seven mtDNA coded subunits (ND1,2,3,4,4L,5,6); Complex III (ubiquinol:cytochrome c oxidoreductase) has one mtDNA coded subunit (cytochrome b); Complex IV (cytochrome c oxidase) has three mtDNA coded subunits (COI, COII, COIII); and Complex V (ATP synthase) has two mtDNA coded polypeptides (subunits 6 and 8). The mtDNA also codes for a complete set of tRNAs and the 12S and 16S rRNAs necessary for mitochondrial protein synthesis^{11,12,23}.

The cytoplasmic location, high copy number, and elevated mutation rate of the mtDNA result in a unique mitochondrial genetics^{11,12,23}. The mtDNA is maternally inherited^{26,27}, and intracellular mixtures of mutant and normal mtDNAs (heteroplasmy) segregate during mitotic and meiotic replication²⁸⁻³¹. Different tissues and organs rely on mitochondrial OXPHOS to different extents. Therefore, as OXPHOS declines because of increasingly severe mtDNA mutations, organ specific energetic thresholds are traversed, yielding variable clinical phenotypes^{31,32,33}. OXPHOS enzyme activities decline with age³⁴⁻³⁶, concomitantly with the age-related accumulation of mtDNA damage in stable tissues³⁷⁻⁴⁰. This may accentuate inherited OXPHOS defects as individuals age, leading to clinical manifestations late in life^{11,12}.

As a result of the quantitative aspects of mtDNA genetics, families harboring deleterious mtDNA mutations frequently show highly variable phenotypic expression among maternal relatives^{11,12}. This is seen for both heteroplasmic mtDNA mutations which segregate between

generations^{31,33} and homoplasmic mutations⁴¹⁻⁴³, whose expression can be influenced by additional mtDNA mutations^{42,44-46}, nuclear DNA alleles⁴⁷, and/or environmental factors. OXPHOS diseases are further complicated by the fact that different mtDNA mutations can act alone or in synergistic groups to produce the same phenotype⁴². This is best exhibited in Leber's hereditary optic neuropathy (LHON), where the same disease has been associated with mutations in the Complex I ND1, ND2, ND4, and ND5 genes^{41,42,44,45,48-50}, in the Complex III cytochrome b gene^{42,46}, and in the Complex IV COI gene⁵¹.

MtDNA mutations which cause LHON fall into two categories: (1) nucleotide substitutions with low pathogenicity that exist as rare polymorphisms in the general population and increase the probability of expressing the disease phenotype^{42,44-46} and (2) nucleotide substitutions with high pathogenicity that cause maternally transmitted disease^{41,48-50}. The low pathogenicity mutations typically cause sporadic disease within families and can be identified through phylogenetic analysis by the clustering of patients around specific mtDNA haplotypes^{42,44,51}. Within these pathological mtDNA lineages, sequential mutations frequently accumulate, increasing the probability of clinical manifestations^{42,44}. The highly pathogenic mutations frequently cause familial disease and occur randomly in the population in association with a variety of mtDNA haplotypes⁵².

Moreover, because of the continued importance of Alzheimer's disease and Parkinson's disease, there is an urgent need to diagnose as well as predict a predisposition to the diseases. This invention satisfies this need by demonstrating that defects in mtDNA are associated with Alzheimer's disease and Parkinson's disease and can be used to predict the likelihood of developing Alzheimer's disease and Parkinson's disease and used to diagnose Alzheimer's disease and Parkinson's disease.

SUMMARY OF THE INVENTION

5 This invention provides a method of diagnosing or predicting a predisposition to Alzheimer's disease and/or Parkinson's disease. The method comprises detecting in a sample from a subject the presence of a mutation, for example, in nucleotide position 4,336, 3,397, 3,196 or an insertion between positions 956 and 965, of mtDNA. The presence of the mutation indicates the presence of or a predisposition to disease. Since each mutation increases the likelihood of developing or having Alzheimer's disease, the detection of more than one of the mutations in an individual can increase the probability of having or developing the disease.

15 The invention also provides a method of determining mutations associated with the presence of or predisposition to Alzheimer's disease and/or Parkinson's disease. The method comprises:

- (a) obtaining a mtDNA-containing sample from a subject with Alzheimer's disease;
- (b) determining the presence of mutations in the mtDNA;
- 20 (c) comparing the mutations to mutations found in a normal subject; and
- (d) determining which mutations have a greater rate of occurrence in the subject with Alzheimer's disease.

25

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a phylogenetic tree generated by parsimony analysis showing the evolutionary relationship between 71 AD, AD+PD, and PD cases, 74 Caucasian controls, and the Cambridge (CA) sequence⁵⁷. This tree is 164 mutational steps in length with a consistency index of 0.756 and represents one of 100 very similar trees of the same length generated by the Tree-Bisection-Reconnection (TBR) branch swapping method of PAUP. The horizontal branch lengths are proportional to the number of mutational events which separate haplotypes. Italicized numbers on top of branches indicate nucleotide positions of mutations discussed in the text. The mtDNA^{16,303-4,336} lineage is highlighted by bold lines. Two additional patients, AD10 and ADPD2, who are outside this lineage but discussed in the text, are designated by boxes.

Polymorphic sites found in all individuals are indicated below by nucleotide position, diagnostic restriction enzyme, and individual code number. Sites are numbered from the first nucleotide of the recognition sequence. A plus (+) indicates a site gain and a minus (-) indicates a site loss relative to the published sequence⁵⁷. Diagonal lines separating restriction sites indicate that a single mutation alters the recognition sequence of more than one enzyme. These sites were considered to be a single site polymorphism in the statistical and phylogenetic analyses. All samples analyzed differ from the published sequence for the following restriction sites: 13702- [HaeIII], 14199- [HincII], and 14268+ [HinfI]. The complete haplotype data set is as follows: 64+ [DdeI] C12; 255+ [HhaI] C21, AD12; 259+ [AluI] C4; 717+ [HinfI] ADPD7; 951- [MboI] AD7, AD10; 956 to 965 [5 bp insertion] ADPD29; 1004- [HincII] ADPD5; 1715- [DdeI] C11, C22, C23, C29, C36, C51, C71, ADPD12, ADPD17, ADPD24, AD29; 1718+ [AluI] C18-C20, ADPD2, ADPD3, AD11, AD17; 1923- [DdeI] C43; 1941+ [HhaI] C49; 2293+ [AluI] AD6; 3192- [DdeI] AD10; 3391+ [HaeIII] AD16, AD32; 3397+ [RsaI] ADPD1, ADPD2; 3759+ [HincII] C51; 4026- [MboI] C56; 4310- [AluI] ADPD8; 4332+ [AvaII] C48, ADPD1, ADPD7, ADPD28, ADPD30, AD23, AD33, PD7, PD8; 4360- [HinfI] C41; 4464- [RsaI] ADPD16; 4529- [HaeII] C11, C23, C29, C49, C71, ADPD12, ADPD17, ADPD24, AD29; 4643+

[RsaI] C7, ADPD20, AD3, AD22, PD1; 4685- [AluI] AD20; 4732/35+ [RsaI]
 C20, ADPD2, AD11; 4745+ [RsaI] ADPD10; 4769- [AluI] all except AD7,
 AD10; 4793+ [HaeIII] C28, C55, ADPD6; 5003- [DdeI] C25; 5198+ [HinfI]
 C1, C57; 5261- [HaeIII] C42, ADPD5; 5584- [AluI] C11, C56; 5823-
 5 [AluI] C15-C17, C25, C56; 6260- [HaeIII] C21, AD13; 6296- [DdeI]
 ADPD6, ADPD19; 6377- [DdeI] AD30; 7025+ [AluI] C1-C27, C29, C32, C34,
 C36, C37, C41, C42, C50-C52, C57, C58, C64, C66, C68, C69, C71, C72,
 ADPD2, ADPD3, ADPD5, ADPD9, ADPD11, ADPD12, ADPD14, ADPD17, ADPD18,
 ADPD20, ADPD22-ADPD25, AD1, AD3-AD6, AD8, AD9, AD11, AD13-AD17, AD19,
 10 AD22, AD24, AD27, AD29-AD32, PD1-PD6; 7055- [AluI] AD30; 7474- [AluI]
 C27, C32; 7570+ [MboI] C36; 7697/7702+ [RsaI] C7, ADPD20, AD3, AD22;
 8012- [RsaI] C52; 8165+ [HaeIII] C51; 8249+ [AvaII]/8250- [HaeIII]
 C11, C23, C29, C41, C51, C68, C71, ADPD12, ADPD17, ADPD24, AD29; 8391+
 15 [AvaII]/8391- [HaeIII] C22, C36; 8515- [DdeI] ADPD24; 8616- [MboI] C29,
 C52, C71, ADPD12, AD29; 8678+ [AluI] ADPD23, AD5; 8858+ [HhaI] all
 except C8, C12; 8894- [HaeIII] C6, C41, C68; 9025- [HaeIII] ADPD16;
 9052-[HaeII]/9053- [HhaI] C9, C12, C21, C24, C58, C69, ADPD23, ADPD25,
 AD5, AD8, AD15, AD23, AD31, PD2-PD5; 9147+ [DdeI] C19; 9253+ [HaeIII]
 C49; 9266- [HaeIII] AD6; 9299+ [AluI] C67; 9380- [HhaI] C39, C49, C61,
 20 AD25; 9714+ [HaeIII] ADPD23, AD5, PD2; 9984+ [HinfI] C50, AD19; 10028+
 [AluI] C11, C23, C29, C71, ADPD12, ADPD17, ADPD24, AD29; 10084+ [TaqI]
 ADPD18; 10097+ [HaeIII] C41; 10394+ [DdeI] C1, C9, C11, C13, C14, C21,
 C23, C24, C27, C29, C32, C37, C52, C57, C58, C64, C69, C71, ADPD11,
 ADPD12, ADPD14, ADPD17, ADPD18, ADPD24, ADPD25, AD4, AD8, AD14-AD16,
 25 AD29, AD31, AD32, PD3-PD5; 10407+ [RsaI] C41; 10598- [AluI] C18, C57;
 10631- [DdeI] C54; 10934+ [MboI] C61; 10971- [HinfI] C50; 11001+
 [HaeII]/11002+ [HhaI] ADPD18; 11313+ [AluI] C7, AD3, PD1; 11362-
 [AluI] C51; 11425+ [AluI] C54; 11439+ [MboI] C72; 11469+ [AluI]
 ADPD25, AD8, PD3; 12345+ [RsaI] C31, C50; 12560- [AluI] C18, ADPD11;
 30 12763+ [AluI] C15; 12810+ [RsaI] C66; 12946+ [DdeI]/12949+
 [HaeII]/12950+ [HhaI] C7; 13018+ [HaeIII] ADPD8, AD18; 13325- [RsaI]
 C5, C7; 13366+ [BamHI]/13367- [AvaII]/13367+ [MboI] C8, C10, C72,
 ADPD2, AD27; 13404- [TaqI] C59; 13635+ [TaqI] C18, C19, C20, ADPD2,
 ADPD3, AD11, AD17; 13957- [HaeIII] C18, ADPD6; 14139+ [MspI] AD6;
 35 14304- [AluI] ADPD23, AD5, PD2; 14347+ [RsaI] C25; 14368- [HinfI] all
 except ADPD10; 14858- [HaeII]/14859- [HhaI] C30; 14869- [MboI] ADPD10,
 AD21; 14899+ [AluI] ADPD15; 15346+ [RsaI] C26; 15434+ [DdeI] C35;

15606+ [AluI] C10, C30, C72, AD27; 15754+ [DdeI] C11, ADPD17, AD1;
 15812- [RsaI] C32; 15861+ [TaqI] C28; 15872+ [RsaI] C50; 15883-
 [HaeIII] ADPD26; 15894/97+ [RsaI] PD5; 15907+ [RsaI] AD19; 15925-
 [MspI] C10, C22, C36, C72, AD27; 16049- [RsaI] AD19; 16065- [HinfI]
 5 C27, C32, C37, C57, C64, ADPD11, ADPD14, AD4, AD14, AD16, AD32; 16145+
 [HaeIII] AD30; 16180+ [MboI] C51; 16208- [RsaI] C31, C63; 16216+
 [HincII] C66; 16238+ [TaqI] C74; 16303- [RsaI] C34, C48, ADPD1, ADPD7,
 ADPD28, ADPD29, ADPD30, AD23, AD33, PD7, PD8; 16310- [RsaI] AD23,
 ADPD28; 16389+ [BamHI]/16390+ [MboI]/16390- [AvaII] C11, C18, C23,
 10 C29, C71, ADPD12, ADPD17, ADPD24, AD29; 16389+ [HinfI]/16390- [AvaII]
 C51, AD6; 16398+ [HaeIII] C26, PD6, ADPD22, AD13; 16436+ [RsaI] C5;
 16467/78+ [DdeI] C39, C61, AD25; 16494+ [MspI] C9; 16517+ [HaeIII] C7,
 C9-C11, C13-C17, C21-C24, C28, C30-C33, C35, C36, C38, C40-C47, C50-
 C56, C58-C60, C62, C63, C65-C74, ADPD4, ADPD6, ADPD8, ADPD11-ADPD13,
 15 ADPD15-ADPD21, ADPD23-ADPD27, AD2, AD3, AD5, AD6, AD8, AD12, AD15,
 AD18-AD22, AD26-AD29, AD31, PD1-PD5, PD8.

The haplotype of the Senegalese used as the African outgroup was
 907+ [TaqI], 2390+ [MboI], 2758- [RsaI], 3592+ [HpaI], 4769- [AluI],
 20 7025+ [AluI], 7055- [AluI], 8858+ [HhaI], 10394+ [DdeI], 10806+
 [HinfI], 13702- [HaeIII], 14199- [HincII], 14268+ [HinfI], 14368-
 [HinfI], 16517+ [HaeIII].

Figure 2 shows the nucleotide sequence of the np^{4,336} mutation
 25 from a control (C) and patient (P). Dideoxy nucleotide sequencing
 reactions for A, C, G, and T are loaded left to right. The A to G
 transition was sequenced on the complementary strand and is indicated
 by arrows as a T to C transition.

Figure 3 shows the AvaII digestion of a 1142 np PCR segment
 30 encompassing the tRNA^{Gln4,336} mutation. Lane 1 is a control in which the
 fragment remains undigested. Lanes 2 to 11 are the nine AD, AD+PD,
 and PD patients identified in the mtDNA^{16,304-4,336} lineage (Figure 1).
 The tRNA^{Gln4,336} mutation causes the 1142 np fragment to be cut into 735
 35 and 407 np fragments.

Figure 4 shows the secondary structure of the tRNA^{Gln} showing the position of the np 4336 point mutation (shaded). Standard base pairing: lines or dashes; non-standard base pairing: asterisks. The homology surrounding the np 4,336 mutation is presented for vertebrates, with the nucleotides corresponding to the np 4,336 mutation indicated by an asterisk. Nucleotides identical with the human sequence are shaded.

Figure 5 shows the nucleotide sequence of the np 3,397 mutation from a control (C) and patient ADPD1 (P). Dideoxy nucleotide sequencing reactions for A, C, G, and T are loaded left to right. The A to G transition at np 3397 is indicated by arrows.

Figure 6 shows the RsaI digestion of a 730 np PCR fragment encompassing the ND1 np 3,397 mutation. Lane 1 is a control in which the 390 np fragment remains undigested. Lanes 2 and 3 are patients ADPD1 and ADPD2 in which the fragment is cut into 330 and 60 np fragments.

Figure 7 shows the evolutionary conservation of methionine 31 in the human ND1 polypeptide altered by the ND1^{3,397 [Met→Val]} mutation. The Polymera and Oligomera are the two developmental lineages of animals. The adjacent tyrosine at amino acid 30 has been proposed to be a cause of LHON⁴⁴. Shaded regions are conserved relative to the human sequence. A, alanine; F, phenylalanine; G, glycine; H, histidine; I, isoleucine; L, leucine; M, methionine; N, asparagine; Q, glutamine; R, arginine; S, serine; V, valine; Y, tyrosine. Oligomera sequences: 1⁵⁷, 2⁸⁸, 3⁸⁹, 4⁹⁰, 5⁹¹, 6⁹², 7, 893. Polymera sequences: 1⁹⁴, 2, 3⁹⁵. Plant sequence: 1⁹⁶. Fungi sequences: 1, 2⁵⁶, 3⁹⁷. Protist sequences: 1⁹⁸, 2⁵⁶, 3⁹⁶.

Figure 8 shows the D-loop sequence analysis of the two AD+PD cases with the ND1^{3,397 [Met→Val]} mutation. Arrows designate sequence divergence of ADPD1 and ADPD2 with the standard Cambridge sequence. The nucleotides at np 4,336 in tRNA^{Gln} and np 3,397 in ND1 are also indicated.

Figure 9 shows the AluI, DdeI, and MboI digestion of a 582 np PCR fragment encompassing the 12S^{956-965ins} mutation. Lanes 1 and 3 are controls showing the normal length fragment. Lane 2 ADPD29 showing approximately a five np increase in length in the normal fragments for AluI (115→120 nps), DdeI (131→136 nps), and MboI (211→216 nps)

Figure 10 shows the proposed secondary structure of the 12S rRNA gene⁹⁹ surrounding the 12S^{956-965ins} insertion. The region of the insertion was determined by direct nucleotide sequencing on each strand. The precise location of the inserted Cs could not be unambiguously determined, but was localized between nps 956 and 965 (bracketed region and shaded nucleotides). Standard base pairing: lines or dashes; non-standard base pairing: asterisks.

Figure 11 shows the restriction endonuclease digests of 16S rRNA PCR fragments that identify the 16S^{3,192} and 16S^{3,196} mutations in patients and controls. Load order for both gels: Lane 1 is a control; 2 is the American Indian with the 16S^{3,192} mutation; 3 is patient AD10 who is heteroplasmic for the 16S^{3,196} mutation, and 4 is an Asian with the 16S^{3,192} mutation. DdeI digests in which the presence of either the 16S^{3,192} or 16S^{3,196} mutation results in a site loss, changing the 342 and 194 np fragments to a 426 np fragment. SspI digests distinguish between the 16S^{3,192} and 16S^{3,196} mutations. A 617 np segment is cut into 530 and 87 np fragments only when the G to A 16S^{3,196} mutation is present. Both the DdeI and SspI digests demonstrate the heteroplasmy in AD10 (Lane 3, both gels). The bottom panel details the relationship of the 16S^{3,192} and 16S^{3,196} mutations to the recognition sequences for the restriction enzymes.

Figure 12 shows the proposed secondary structure of the 16S rRNA at the 3'-end¹⁰⁰ in the region of the 16S^{3,196} mutation. The polymorphic nucleotides at nps 3,192 and 3,196 are indicated by shaded boxes. Standard base pairing: lines or dashes; non-standard base pairing: asterisk.

Figure 13 shows the restriction endonuclease digests identifying the G to A ND2⁵⁴⁶⁰ mutation in patients and controls. Left panel, HphI

cuts the 332 np fragment to 198 and 134 np fragments in individuals with an A at np 5460. Right panel, *AccI* digest this fragment generated by a modified PCR primer only when there is a G at np 5460, cutting the 208 np fragment to 190 and 18 np fragments. Any

5 nucleotide change at np 5460 eliminates this site. Load order for both panels: Lanes 2 to 5 have the Cambridge sequence⁵⁷, while lanes 1, 6, and 7 harbor the G to A mutation.

DETAILED DESCRIPTION OF THE INVENTION

5 This invention provides a method of diagnosing or predicting a predisposition to Alzheimer's disease and/or Parkinson's Disease. The method comprises detecting in a sample from a subject the presence of a mutation, for example, in nucleotide positions 4,336, 3,397, 3,196 or an insertion between positions 956 and 965, of mtDNA. The presence of the mutation indicates the presence of or a predisposition to Alzheimer's disease and/or Parkinson's disease. Since each mutation increases the likelihood of developing or having Alzheimer's/Parkinson's disease, the detection of more than one of the mutations in an individual can increase the probability of having or developing the disease.

15 The invention also provides a general method of determining mutations associated with the presence of or predisposition to Alzheimer's disease and/or Parkinson's disease. The method comprises:

- (a) obtaining a mtDNA-containing sample from a subject with Alzheimer's disease;
- 20 (b) determining the presence of mutations in the mtDNA;
- (c) comparing the mutations to mutations found in a normal subject; and
- (d) determining which mutations have a greater rate of occurrence in the subject with Alzheimer's/Parkinson's disease.

25 Since the clinical and neuropathological features of Alzheimer's disease and Parkinson's disease overlap extensively, the mutations can contribute to either or both diseases. Therefore, the diseases may sometimes simply be referred to herein as Alzheimer's/Parkinson's disease.

30 As used herein, "predicting a predisposition to Alzheimer's and/or Parkinson's disease" means determining an increased probability of developing Alzheimer's/Parkinson's disease during the subject's lifetime. Depending on the type and number of mutations correlated with the disease, it is possible to predict the likelihood of an individual developing disease. By "correlated" is meant a mutation

35

which occurs in a higher rate in Alzheimer's/Parkinson's subjects than in normal subjects.

As used herein, "diagnosing" means using the presence of a mutation associated with Alzheimer's/Parkinson's disease as a factor in making an Alzheimer's/Parkinson's disease diagnosis. Since the presence of such mutations does not mean the subject has Alzheimer's/Parkinson's disease, the detection of the mutation is merely a step in the disease state diagnosis.

As used herein, "isolated" means free of at least some of the contaminants associated with the nucleic acid or polypeptides occurring in a natural environment.

As used herein, "nucleic acid characteristic of human mitochondrial DNA" refers to a nucleic acid which has sufficient nucleotides surrounding the codons at the mutation positions to distinguish the nucleic acid from nucleic acids encoding non-related genes. The specific length of the nucleic acid is a matter of routine choice based on the desired function of the sequence. For example, if one is making probes to detect the mutation in nt position 4,336, the length of the nucleic acid is preferably small, but must be long enough to prevent hybridization to undesired background sequences. However, if the desired hybridization is to a nucleic acid which has been amplified, background hybridization is less of a concern and a smaller probe can be used. In general, such a probe will be between 10 and 100 nucleotides, especially between 10 and 40 and preferably between 15 and 25 nucleotides in length. It is apparent to one of skill in the art that nucleotide substitutions, deletions, and additions may be incorporated into the polynucleotides of the invention. However, such nucleotide substitutions, deletions, and additions should not substantially disrupt the ability of the polynucleotide to hybridize under conditions that are sufficiently stringent to result in specific hybridization.

As used herein, the term "mutant" generally refers to a mutation in the mtDNA associated with Alzheimer's disease and/or Parkinson's

disease. "Mutant" can specifically refer to a mutation at nucleotide position 4,336, 3,397, 3,196 or an insertion between positions 956 and 965 of mtDNA. For mutations at nucleotide position 4,336, 3,197 or the insertion between 956 to 965, a structural RNA is altered which will affect mitochondrial gene expression. For the mutation at nucleotide position 3,397, a codon is changed from specifying methionine to valine. Thus, other mutations which cause a change in the codon from methionine can be associated with Alzheimer's/Parkinson's disease, especially where the change is to valine.

"Specific or selective hybridization" as used herein means the formation of hybrids between a probe nucleic acid (e.g., a nucleic acid which may include substitutions, deletions, and/or additions) and a specific target nucleic acid (e.g., a nucleic acid having the mutated sequence), wherein the probe preferentially hybridizes to the specific target such that, for example, a band corresponding to the mutated DNA or restriction fragment thereof can be identified on a Southern blot, whereas a corresponding normal or wild-type DNA is not identified or can be discriminated from a variant DNA on the basis of signal intensity. Hybridization probes capable of specific hybridization to detect a single-base mismatch may be designed according to methods known in the art^{124, 125, 126, 127, 128}.

The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials such as bacteria, plants, fungi, or animal (particularly mammalian) cells or tissues. Agents are evaluated for potential biological activity by inclusion in screening assays.

As discussed below in greater detail, the mutation can be detected by many methods. For example, the detecting step can comprise combining a nucleotide probe capable of selectively hybridizing to a nucleic acid containing the mutation with a nucleic acid in the sample and detecting the presence of hybridization. Additionally, the detecting step can comprise amplifying the

nucleotides of the mutation and detecting the presence of the mutation in the amplified product. Further, the detecting step can comprise selectively amplifying the nucleotides of the mutation and detecting the presence of amplification. Finally, the detecting step can
5 comprise detecting the loss or gain of a restriction fragment created by an enzyme digest of the nucleotides of the mutation.

Detection Techniques

10 Once the location of the mutations are known and associated with Alzheimer's/Parkinson's disease, the methods to detect the mutations are standard in the art. The sequence of various nucleotide probes can be determined from the known sequence of mtDNA¹²⁹, especially the sequences surrounding the mutations and insertions.

15 Detection of point mutations using direct probing involves the use of oligonucleotide probes which may be prepared, for example, synthetically or by nick translation. The probes may be suitably labeled using, for example, a radio label, enzyme label, fluorescent
20 label, biotin-avidin label and the like for subsequent visualization in the example of Southern blot hybridization procedure. The labeled probe is reacted with a bound sample DNA, e.g., to a nitrocellulose sheet under conditions such that only fully complementary sequences hybridize. The areas that carry DNA sequences complementary to the
25 labeled DNA probe become labeled themselves as a consequence of the reannealing reaction. The areas of the filter that exhibit such labeling may then be visualized, for example, by autoradiography. The labeled probe is reacted with a DNA sample bound to, for example, nitrocellulose under conditions such that only fully complementary
30 sequences will hybridize. Tetra-alkyl ammonium salts bind selectively to A-T base pairs, thus displacing the dissociation equilibrium and raising the melting temperature. At 3M Me 4NCl this is sufficient to shift the melting temperature to that of G-C pairs. This results in a marked sharpening of the melting profile. The stringency of
35 hybridization is usually 5°C below the T_i (the irreversible melting temperature of the hybrid formed between the probe and its target sequence) for the given chain length. For 20mers the recommended

hybridization temperature is about 58°C. The washing temperatures are unique to the sequence under investigation and need to be optimized for each variant.

5 Alternative probing techniques, such as ligase chain reaction (LCR), involve the use of mismatch probes, i.e., probes which are fully complementary with the target except at the point of the mutation. The target sequence is then allowed to hybridize both with
10 oligonucleotides which are fully complementary and have oligonucleotides containing a mismatch, under conditions which will distinguish between the two. By manipulating the reaction conditions, it is possible to obtain hybridization only where there is full complementarity. If a mismatch is present there is significantly reduced hybridization.

15 The polymerase chain reaction (PCR) is a technique that amplifies specific DNA sequences with remarkable efficiency. Repeated cycles of denaturation, primer annealing and extension carried out with polymerase, e.g., a heat stable enzyme Taq polymerase, leads to
20 exponential increases in the concentration of desired DNA sequences. Given a knowledge of the nucleotide sequence of the mutations, synthetic oligonucleotides can be prepared which are complementary to sequences which flank the DNA of interest. Each oligonucleotide is complementary to one of the two strands. The DNA can be denatured at
25 high temperatures (e.g., 95°C) and then reannealed in the presence of a large molar excess of oligonucleotides. The oligonucleotides, oriented with their 3' ends pointing towards each other, hybridize to opposite strands of the target sequence and prime enzymatic extension along the nucleic acid template in the presence of the four
30 deoxyribonucleotide triphosphates. The end product is then denatured again for another cycle. After this three-step cycle has been repeated several times, amplification of a DNA segment by more than one million-fold can be achieved. The resulting DNA may then be directly sequenced in order to locate any genetic alteration.
35 Alternatively, it may be possible to prepare oligonucleotides that will only bind to altered DNA, so that PCR will only result in multiplication of the DNA if the mutation is present. Following PCR,

direct visualization or allele-specific oligonucleotide hybridization¹³⁰ may be used to detect the Alzheimer's/Parkinson's disease point mutation. Alternatively, an adaptation of PCR called amplification of specific alleles (PASA) can be employed; this uses differential amplification for rapid and reliable distinction between alleles that differ at a single base pair. Other techniques, such as 3SR, which utilize RNA polymerase to achieve high copy number, can also be used where appropriate.

In yet another method, PCR may be followed by restriction endonuclease digestion with subsequent analysis of the resultant products. As shown in the examples, the substitution of G for A at base pair 4,336, results in a gain of an *Ava*II site. The gain of this restriction endonuclease recognition site facilitates the detection of the Alzheimer's/Parkinson's disease mutation using restriction fragment length polymorphism (RFLP) analysis or by detection of the presence or absence of a polymorphic *Ava*II site in a PCR product that spans base pair 4,336.

For RFLP analysis, DNA is obtained, for example from the blood of the subject suspected of having Alzheimer's/Parkinson's disease and from a normal subject, is digested with a restriction endonuclease, and subsequently separated on the basis of size by agarose gel electrophoresis. The Southern blot technique can then be used to detect, by hybridization with labeled probes, the products of endonuclease digestion. The patterns obtained from the Southern blot can then be compared. Using such an approach, DNA spanning an Alzheimer's/Parkinson's mutation, such as an *Ava*II site, is detected by determining the number of bands detected and comparing this number to the normal subject. Various restriction endonucleases can be utilized effectively for each mutation.

Similar creation of additional restriction sites by nucleotide substitutions at the disclosed mutation sites can be readily calculated by reference to the genetic code and a list of nucleotide sequences recognized by restriction endonucleases¹³¹.

Single strand conformational analysis (SSCA)^{132,133} offers a relatively quick method of detecting sequence changes which may be appropriate in at least some instances.

5 In general, primers for PCR and LCR are usually about 20 bp in length and the preferable range is from 15-25 bp. Better amplification is obtained when both primers are the same length and with roughly the same nucleotide composition. Denaturation of strands usually takes place at 94°C and extension from the primers is usually
10 at 72°C. The annealing temperature varies according to the sequence under investigation. Examples of reaction times are: 20 mins denaturing; 35 cycles of 2 min, 1 min, 1 min for annealing, extension and denaturation; and finally a 5 min extension step.

15 PCR amplification of specific alleles (PASA) is a rapid method of detecting single-base mutations or polymorphisms^{134,135,136,137,138,139,140}. PASA (also known as allele specific amplification) involves amplification with two oligonucleotide primers such that one is
20 allele-specific. The desired allele is efficiently amplified, while the other allele(s) is poorly amplified because it mismatches with a base at or near the 3' end of the allele-specific primer. Thus, PASA or the related method of PAMSA may be used to specifically amplify the mutation sequences of the invention. Where such amplification is done
25 on genetic material (or RNA) obtained from an individual, it can serve as a method of detecting the presence of the mutations.

 As mentioned above, a method known as ligase chain reaction (LCR) can be used to successfully detect a single-base
30 substitution^{150,151}. LCR probes may be combined or multiplexed for simultaneously screening for multiple different mutations. Thus, LCR can be particularly useful where, as here, multiple mutations are predictive of the same disease.

35

Transgenic Animals and Cell Lines and Screening Methods

5 Having identified subjects having Alzheimer's/Parkinson's disease associated with point and insertion mutations, the subjects can be screened for an agent capable of treating Alzheimer's and/or Parkinson's disease. This method comprises administering the agent to an animal, including a human, having a mutation in mtDNA associated with Alzheimer's/Parkinson's disease and monitoring the effect of the agent on the animal's condition. If the symptoms of Alzheimer's and/or Parkinson's disease improve, the agent can be used as a treatment for the disease.

15 In addition, it is possible, using genetic manipulation, to develop transgenic model systems and/or whole cell systems containing the mutations for use, for example, as model systems for screening for drugs and evaluating drug effectiveness. Additionally, such model systems provide a tool for defining the underlying biochemistry of mtDNA, which thereby provides a basis for rational drug design.

20 One type of cell system can be naturally derived. For this, blood samples from an affected subject must be obtained in order to provide the necessary cells which can be permanently transformed into a lymphoblastoid cell line using, for example, Epstein-Barr virus. Once established, such cell lines can be grown continuously in suspension culture and may be used for a variety of in vitro experiments to study expression and processing. An alternative method for constructing a cell line is to genetically engineer the mutated gene into an established cell line of choice.

30 In yet a further use of the present invention, the disease mutation can be introduced into transgenic animals. For example, a mutated gene can be isolated by cloning (e.g., λ Charon35, cosmid, retrovirus or yeast artificial chromosome). The mtDNA genetic code would be altered by in vitro mutagenesis to create a mRNA compatible with the nuclear-cytosol system. A mitochondrial targeting sequence would then be added to the amino terminal end by gene splicing from a nuclear OXPHOS gene such as the B subunit of the ATP synthase. The

35

chimeric gene could be inserted in an appropriate nuclear vector. Alternatively, the gene could be inserted into a mitochondria-specific vector. The vector could be transferred or inserted into the cell or animal and the products screened for appropriate integration. As a result of the transfer, the resultant transgenic non-human animal will express the Alzheimer's/Parkinson's genes.

One approach to creating transgenic animals is to mutate the animal mtDNAs by in vivo mutagenesis, transfer the mutant mtDNA into the mitochondrial embryonic stem cells by DNA transfection and injection of the embryonic stem cells into blastocysts to prepare allophenic mice carrying the disease causing mtDNA mutation¹⁵². Alternatively, the technique of microinjection of the mutated gene or mitochondria, into a one-cell embryo followed by incubation in a foster mother can be used. Certain possibilities for the general use of transgenic animals are disclosed^{153,154,155,156,157}. Alternatively, viral vectors, e.g., Adeno-associated virus, can be used to deliver the mutated gene to the stem cell. In addition, such viral vectors can be used to deliver the mutated gene to a developed animal and then used to screen^{158,159}.

To more accurately model the disease, transgenic animals can be created with a combination of multiple mutations, e.g. 4,336, 3,397, 3,196 or the insertion mutation, in the same animal produced by the technologies described above can be utilized. Similarly, cell lines manipulated to include several mutations will be of value in modeling the disease and are an important consequence of this invention.

Gene Therapy

Having detected the genetic mutation in the mitochondrial gene sequence in an individual, it is possible to employ gene therapy, in the form of gene implants, to prevent the development of the disease. In the present case, if significant mutations in the mtDNA are discovered, the nucleus from the egg of a carrier mother can be inserted into an egg having normal mtDNA from which the original nucleus has been removed. The egg can then be reintroduced into the

mother. Such a method can be utilized to delete any mutation of the mtDNA while maintaining the genes from the mother which affect appearance.

5 Antibodies and Recombinant Expression of Mutant Polypeptides

When the mutated oxphos gene product is a polypeptide, e.g. the 3,397 mutation, it can be used to prepare antisera and monoclonal antibodies using, for example, the method of Kohler and Milstein¹⁶⁰.
10 Such monoclonal antibodies could then form the basis of a diagnostic test.

Such variant polypeptides can also be used to immunize an animal for the production of polyclonal antiserum¹⁶¹. For example, a
15 recombinantly produced fragment of a variant polypeptide can be injected into a mouse along with an adjuvant so as to generate an immune response. Murine immunoglobulins which bind the recombinant fragment with a binding affinity of at least 1×10^7 m-1 can be harvested from the immunized mouse as an antiserum, and may be further
20 purified by affinity chromatography or other means. Additionally, spleen cells are harvested from the mouse and fused to myeloma cells to produce a bank of antibody-secreting hybridoma cells. The bank of hybridomas can be screened for clones that secrete immunoglobulins which bind the recombinantly produced fragment with an affinity of at
25 least 1×10^6 M-1. More specifically, immunoglobulins that selectively bind to the variant polypeptides but poorly or not at all to wild-type polypeptides are selected, either by pre-absorption with wild-type proteins or by screening of hybridoma cell lines for specific idiotypes that bind the variant but not wild-type.

30

The nucleic acid sequences of the present invention capable of ultimately expressing the desired variant polypeptides can be formed from a variety of different polynucleotides (genomic or cDNA, RNA, synthetic oligonucleotides, etc.) as well as by a variety of different
35 techniques.

The DNA sequences can be expressed in hosts after the sequences have been operably linked to, i.e., positioned to ensure the functioning of, an expression control sequence. These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors can contain selection markers, e.g., tetracycline resistance or hygromycin resistance, to permit detection and/or selection of those cells transformed with the desired DNA sequences (see, e.g., U.S. Patent 4,704,362).

Polynucleotides encoding a variant polypeptide may include sequences that facilitate transcription (expression sequences) and translation of the coding sequences such that the encoded polypeptide product is produced. Construction of such polynucleotides is well known in the art¹²⁴. For example, such polynucleotides can include a promoter, a transcription termination site (polyadenylation site in eukaryotic expression hosts), a ribosome binding site, and, optionally, an enhancer for use in eukaryotic expression hosts, and, optionally, sequences necessary for replication of a vector.

E. coli is one prokaryotic host useful particularly for cloning DNA sequences of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In these prokaryotic hosts one can also make expression vectors, which will typically contain expression control sequences compatible with the host cell (e.g., an origin of replication). In addition, any number of a variety of well-known promoters will be present, such as the lactose promoter system, a tryptophan (Trp) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The promoters will typically control expression, optionally with an operator sequence, and have ribosome binding site sequences for example, for initiating and completing transcription and translation.

Other microbes, such as yeast, may also be used for expression. *Saccharomyces* can be a suitable host, with suitable vectors having

expression control sequences, such as promoters, including 3-phosphoglycerate kinase or other glycolytic enzymes, and an origin of replication, termination sequences etc. as desired.

5 In addition to microorganisms, mammalian tissue cell culture may also be used to express and produce the polypeptides of the present invention¹⁶². Eukaryotic cells are actually preferred, because a number of suitable host cell lines capable of secreting intact human proteins have been developed in the art, and include the CHO cell
10 lines, various COS cell lines, HeLa cells, myeloma cell lines, Jurkat cells, etc. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer¹⁶³, and necessary information processing sites, such as
15 ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferred expression control sequences are promoters derived from immunoglobulin genes, SV40, Adenovirus, Bovine Papilloma Virus, etc. The vectors containing the
20 DNA segments of interest (e.g., polypeptides encoding a variant polypeptide) can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts.

25 Kits

 The method lends itself readily to the formulation of test kits which can be utilized in diagnosis. Such a kit would comprise a carrier compartmentalized to receive in close confinement one or more
30 containers wherein a first container may contain suitably labeled DNA probes. Other containers may contain reagents useful in the localization of the labeled probes, such as enzyme substrates. Still other containers may contain restriction enzymes (such as RsaI), buffers etc., together with instructions for use.

35

Analysis of Mutations

Three lines of evidence suggest that the tRNA^{Gln4,336} mutation defines an mtDNA lineage which is associated with an increased risk for manifesting Alzheimer's disease and Parkinson's disease. First, the tRNA^{Gln4,336} mtDNA lineage correlates with the Alzheimer's disease and Parkinson's disease phenotype, being found 7.4 times more frequently in Caucasian patients than in Caucasian controls. Second, Alzheimer's disease and Parkinson's disease patients cluster in the tRNA^{Gln4,336} lineage, suggesting that these clinical phenotypes are linked to this mtDNA haplotype. Third, the tRNA^{Gln4,336} lineage has been found to harbor two additional novel and potentially pathogenic mutations, the ND1^{3,397[Met→Val]} missense mutation and the 12S^{956-965ins} insertion. Since multiple, sequential, rare events are more likely to cluster together if they are functionally related⁴², the accumulation of these three mtDNA mutations in an Alzheimer's disease and Parkinson's disease mtDNA lineage demonstrates that these mutations act synergistically to increase the risk of manifesting these diseases.

The ND1^{3,397[Met→Val]} mutation has a number of features which demonstrate that it also contributes to Alzheimer's disease and Parkinson's disease pathology. First, the mutation changes the highly conserved methionine at codon 31 in the ND1 polypeptide to a valine. This mutation is immediately adjacent to the ND1^{3,394[Tyr→His]} mutation in codon 30 that has been found in a rare form of LHON⁴⁴. Therefore, mutations in this region of ND1 appear to be pathologically significant. Second, the ND1^{3,397[Met→Val]} mutation is maternally inherited and maternal relatives of pedigrees harboring this mutation have also been found to manifest clinical features of Alzheimer's disease. Third, the mutation has arisen two independent times in Caucasian patients, but has not been found in 248 Caucasian controls, demonstrating that in Caucasians it is associated with AD+PD pathology.

The 12S^{956-965ins} insertion was found in an AD+PD patient (ADPD29) within the mtDNA^{16,304+4,336} lineage. Although small insertions occur

commonly in the non-coding regions of the mtDNA⁵⁵, nucleotide insertion mutations have not been observed within mtDNA genes. This observation was affirmed by the absence of the 12S^{956-965ins} mutation in 699 individuals from various ethnic groups. Thus, while the unique nature of this event precludes calculation of its statistical significance, the location of the insertion within an OXPHOS gene that is essential to protein synthesis, its absence in all controls, and its association with the rare tRNA^{Gln4,336} mutation demonstrates that it can be functionally related to the AD+PD phenotype.

Each of the four mtDNA mutations identified in this study can result in OXPHOS defects similar to those reported in Alzheimer's disease and Parkinson's disease patients. Defects in respiratory Complexes I and IV have been reported for PD^{17,18}, in Complex I for PD^{13-16,18}, and in Complex IV for PD¹⁸ and AD¹⁹. The tRNA^{Gln4,336} and 12S^{956-965ins} mutations would be expected to impair mitochondrial protein synthesis and thus affect Complexes I and IV, as observed for the tRNA^{Lys8,334} mutation that causes MERRF^{31,32}, the tRNA^{Leu3,260} mutation that causes maternally inherited mitochondrial myopathy and cardiomyopathy⁷⁸, and the tRNA^{Leu3,243} mutation that causes MELAS⁷⁴. The ND1^{3,397[Met-Wal]} mutation could cause defects in Complex I as observed for the LHON ND1^{3,460[A1a-Thr]} mutation¹²¹, the ND1^{4,160[Leu-Pro]} mutation¹²³, and the ND4^{11,778[Arg-His]} mutation^{121,122}.

The clustering of the tRNA^{Gln4,336}, the 12S^{956-965ins}, and the ND1^{3,397[Met-Wal]} mutations on the mtDNA^{16,304+4,336} lineage is analogous to the accumulation of multiple mtDNA mutations in LHON^{11,12,42,44}. This demonstrates that several, mildly deleterious mtDNA mutations can accumulate within an mtDNA lineage and act synergistically to cause Alzheimer's disease and Parkinson's disease. The cumulative OXPHOS defect produced by these mutations can be further exacerbated by the age-related decline in OXPHOS function resulting from the accumulation of somatic mtDNA mutations in stable tissues^{11,12}. As the severity of the inherited OXPHOS defect increases, the neuronal ATP generating capacity would decline, increasing the probability of falling below the expression threshold and causing the Alzheimer's disease or Parkinson's disease phenotype^{11,12}.

The probabilistic nature of this model means that individuals who inherit mildly deleterious mtDNA mutations, such as the tRNA^{Gln4,336} mutation, will only occasionally manifest disease symptoms, and appear as seemingly sporadic cases. This is comparable to the situation for the LHON mutations at position 15,257 in the cytochrome b gene and position 7,444 in the COI gene^{42,51}. By contrast, individuals inheriting more severe mtDNA mutations or combinations of mutations such as tRNA^{Gln4,336} plus ND1^{3,397[Met→Val]} who have greater reductions in neuronal ATP generating capacity and thus be more likely to have affected maternal relatives. Other genetic and environmental factors could also influence the neuronal OXPHOS capacity and increase the likelihood of disease expression. Genetic factors could include nuclear genes similar to the X-linked gene proposed to influence LHON expression⁴⁷, and some of the loci affecting Alzheimer's disease and Parkinson's disease might be nuclear encoded OXPHOS genes^{11,12}.

Throughout this application various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

The following examples are intended to illustrate but not limit the invention. While they are typical of those that might be used, other procedures known to those skilled in the art may be alternatively employed.

EXAMPLES

Patient and control samples

The AD plus PD samples included 73 autopsy brains with neuropathological changes of AD+PD⁴, 62 brains with the conventional changes of AD^{5,81} and 38 blood and muscle samples collected from Parkinson's disease patients responsive to levodopa. The age range for these groups of individuals was approximately 51 to 91 years. One brain included in the Alzheimer's disease group which harbored the tRNA^{Gln4,336} mutation was originally considered borderline for

classification in the Alzheimer's disease group based on number of plaques for age. However, further analysis, which included staining of sections with the Alzheimer-50 antibody and silver staining, identified a significant accumulation of neuritic plaques and neurofibrillary tangles and was felt to be consistent with early Alzheimer's disease. Therefore, this case was included in the patient group. However, either exclusion or inclusion of this patient in the analysis did not alter the p-values. All p-values were calculated using the Pearson Chi-square, Yates corrected Chi-square, and Fisher exact (two tail) method⁸².

Additional pathological samples included blood and muscle samples from three patients with multisystems atrophy and one with progressive supranuclear palsy, and brain samples from two patients with amyotrophic lateral sclerosis, one with MELAS, two with multiple sclerosis, one with mental retardation, epilepsy and short stature, and one with Leigh's disease. Control samples included 23 brains with no observed neuropathology, and 547-699 blood and placental samples examined in our laboratory.

Phylogenetic analysis

Genomic DNA was isolated from brains frozen at autopsy, platelets, or skeletal muscle as previously described³⁸. For haplotype analysis, the entire mtDNA genome was amplified as nine overlapping fragments by polymerase chain reaction (PCR) and surveyed for restriction fragment length polymorphisms by digestion with the following 14 restriction endonucleases: AluI, AvaII, BamHI, DdeI, HaeII, HaeIII, HhaI, HinfI, HincII, HpaI, HpaII, MboI, RsaI, TaqI^{18,42,53,54}. These PCR fragments were generated with the following forward → and reverse ← primer pairs and PCR annealing temperature (T_H): (1) nps 1562-1581 →; 3701-3717 ← (T_H=51°C), (2) nps 3007-3023 →, 5898-5917 ← (T_H=55°C), (3) nps 5317-5333 →, 7588-7608 ← (T_H=57°C), (4) nps 7392-7410 →, 8902-8921 ← (T_H=57°C), (5) nps 8282-8305 →, 10,088-10,107 ← (T_H=57°C), (6) nps 9911-9932 →, 11,851-11,873 ← (T_H=69°C), (7) nps 11,673-11,691 →, 13,932-13,950 ← (T_H=57°C), (8) nps 13,914-13,930 →, 16,527-16,547 ← (T_H=47°C), (9) nps 16,453-16,472 →, 1677-1696 ←

($T_H=61^\circ\text{C}$). Restriction fragments were resolved by electrophoresis on 1.0-2.5% NuSieve plus 1.0% SeaKem agarose (FMC BioProducts) gels, visualized by ethidium bromide staining and mapped by the sequence comparison method^{57,83,84}. The composite restriction map which
5 represents individual haplotypes is given in the legend of Figure 1.

The evolutionary relationship among the 71 AD, AD+PD, and PD patients and 74 Caucasian controls plus the Cambridge sequence⁵⁷ was inferred using parsimony analysis with the computer program PAUP⁸⁵. An
10 African mtDNA haplotype from a Senegalese was used to root the resulting phylogenetic tree. Sequence divergence between haplotypes was estimated by a maximum likelihood procedure using the computer program DREST⁸⁶. The divergence rate used to assess restriction site variation was 2-4% per million years⁵⁸. For the D-loop sequence
15 analysis, only those nucleotide positions unambiguously sequenced and shared by the individuals being compared were included in divergence time calculations (900 total nucleotides for the ND1^{3,397[Met-Val]} mutation and 344 total nucleotides for individuals with the tRNA^{Gln4,336} mutation). The divergence rate used to assess differences in the D-loop sequence between individuals was 8.4% per million years⁸⁷.
20

DNA sequencing

D-loop sequencing and sequencing of mutations identified by
25 phylogenetic analysis were performed using previously described direct sequencing protocols^{31,39,42}.

A D-loop PCR segment was generated from primers positioned at either nps 15,705-15,723 → and nps 221-240 ← ($T_H=47^\circ\text{C}$) or nps 15,243-
30 15,259 → and nps 1467-1486 ← ($T_H=45^\circ\text{C}$). Asymmetric PCRs were generated from this fragment for sequencing using the following primer pairs: (1) nps 15,704-15,723 → and 16,344-16,364 ← ($T_H=49^\circ\text{C}$), (2) nps → 5,794-15,813 → and 16,527-16,547 ← ($T_H=51^\circ\text{C}$), (3) nps 16,141-16,160 → and 408-429 ← ($T_H=53^\circ\text{C}$), (4) nps 16,141-16,160 → and 16,401-16,420 ←
35 ($T_H=53^\circ\text{C}$), (5) nps 16,141-16,160 → and 16,527-16,547 ← ($T_H=51^\circ\text{C}$), (6) nps 15,978-15,996 → and 16,527-16,547 ← ($T_H=55^\circ\text{C}$), (7) nps 16,141-16,160 → and 408-429 ← ($T_H=53^\circ\text{C}$), (8) nps 1-20 → and 408-429 ←

($T_H=55^\circ\text{C}$), (9) nps 1-20 \rightarrow and 1227-1246 \leftarrow ($T_H=51^\circ\text{C}$), (10) nps 247-264 \rightarrow and 751-770 \leftarrow ($T_H=51^\circ\text{C}$), (11) nps 242-261 \rightarrow and 751-770 \leftarrow ($T_H=53^\circ\text{C}$).

The following primers were used to sequence the D-loop from the single stranded mtDNA generated by the asymmetric PCR: (1) nps 15,978-15,997, (2) nps 16,028-16,047, (3) nps 16,225-16,244, (4) nps 16,287-16,306, (5) nps 16,364-16,344, (6) nps 16,401-16,421, (7) nps 16,527-16,547, (8) nps 131-150, (9) nps 242-261, (10) nps 371-390, (11) nps 617-636.

The tRNA^{Gln}₃₃₆ mutation was amplified within a 2.9 kb PCR fragment generated from primer pair 2 in the phylogenetic analysis. Single stranded mtDNA was generated using nested primers at nps 3941-3970 \rightarrow and 5049-5067 \leftarrow ($T_H=53^\circ\text{C}$) and sequenced with a primer at nps 4201-4219.

The ND1^{3,397(Met \rightarrow Val)} mutation was sequenced from a 1.4 kb mtDNA fragment amplified with primers at nps 3108-3127 (\rightarrow) and 4489-4508 (\leftarrow) ($T_H=53^\circ\text{C}$). Single stranded mtDNA was generated using primers at nps 3108-3127 (\rightarrow) and 3701-3717 (\leftarrow) ($T_H=51^\circ\text{C}$) and sequenced with a primer at nps 3317-3337.

Attempts to define the sequence of the 12S^{956-965ins} mutation were made using a mtDNA fragment amplified with primers at nps 1-20 \rightarrow and 1696-1677 \leftarrow ($T_H=55^\circ\text{C}$). Numerous asymmetric PCR reactions were prepared and the sequencing primers used were located at nps 811-831, 871-890, and 1064-1045. The 12S956-965ins mutation was localized with this approach to the mtDNA sequence between nps 956 and 965. However, the precise mtDNA sequence within this region which contains nine cytosines in the Cambridge sequence⁵⁷ could not be unambiguously ascertained. The best interpretation of the sequencing gels in this region is an insertion of approximately five cytosines.

The 16S^{3,192} and 16S^{3,196} mutations were sequenced from a 2.9 kb PCR fragment generated from primer pair 2 of the phylogenetic analysis. Single stranded mtDNA was generated using primers at nps 3007-3023 \rightarrow and 3701-3717 \leftarrow ($T_H=51^\circ\text{C}$) and sequenced with a primer at nps 3108-3127.

Restriction endonuclease screening for point mutations

5 Restriction endonuclease screening of patient and control mtDNAs was performed using *Ava*II for the tRNA^{Gln4,336} mutation; *Rsa*I for the ND1^{3,397[Met→Val]} mutation; *Alu*I, *Dde*I, and *Mbo*I for the 12S^{956-965ins} mutation; *Dde*I and *Ssp*I for the 16S^{3,196} mutation; and *Acc*II and *Hph*I for the ND2⁵⁴⁶⁰ mutation. Digestions were performed under standard conditions and separated on either on 2.5-2.8% NuSieve agarose plus 0.8-0.9% SeaKem agarose gels. Fragments were visualized by ethidium bromide
10 fluorescence.

To screen for the *Ava*II site gain produced by the tRNA^{Gln4,336} mutation in patients and controls, a 578 np segment of the mtDNA was amplified at T_m=55°C with primers at nps 3951-3970 plus a 10 np tail → and 4489-4508 plus a 10 np tail ←. The normal sequence remains
15 undigested while those containing the tRNA^{Gln4,336} mutation are cleaved, giving fragments at 395 and 183 nps. A 1142 nps segment of the mtDNA was amplified for Figure 3 using primers at nps 3598-3615 → and 4720-4739 ←. The normal sequence remains undigested giving a fragment of
20 1142 np, while the tRNA^{Gln4,336} mutation yields fragments at 735 and 407 nps.

The *Rsa*I site gain produced by the ND1^{3,397[Met→Val]} mutation was evaluated by amplifying a 730 np segment of the ND1 gene at T_m=50°C
25 with primers at nps 3007-3023 plus a 9 np tail (→) and 3701-3717 plus a 10 np tail (←). Sizes of the digestion products for the normal sequence were 390, 214, and 126 nps and for the mutant containing ND1^{3,397[Met→Val]} were 329, 214, 126, and 61 nps.

30 The 12S^{956-965ins} mutation was detected by multiple restriction endonuclease digestions including *Alu*I, *Dde*I, and *Mbo*I. The 582 np segment of mtDNA containing the 12S^{956-965ins} mutation was amplified at T_m=51°C using primers at nps 664-683 → and 1227-1246 ←. *Alu*I digestion of mtDNA with the Cambridge sequence produced fragments with
35 sizes of 115, 101, 96, 90, 85, 80, and 11 nps. The 12S^{956-965ins} mutation increased the 115 np fragment to approximately 120 nps. *Dde*I digestion produced fragments with sizes of 153, 131, 76, 72, 35, 15, and 9 nps.

The 12S^{956-965ins} mutation increased the 131 np fragment to approximately 136 nps. MboI digestion produced fragments with sizes of 276, 211, 76, and 19 nps. The 12S^{956-965ins} mutation increased the 211 np fragment to approximately 216 nps.

5

The 16S^{3,192} and 16S^{3,196} mutations were evaluated using a 617 np segment of the mtDNA amplified at $T_m=51^\circ\text{C}$ with primers at nps 3108-3127 \rightarrow and 3701-3717 plus a 10 np tail \leftarrow . DdeI digestion resulted in a site loss when either the C to T mutation at np 3,192 or the G to A mutation at np 3,196 were present. Sizes of the digestion products for the normal sequence were 342, 194, and 84 nps and for the sequence containing either the 16S^{3,192} or 16S^{3,196} mutations were 426 and 194 nps. SspI digestion revealed a site gain only when the G to A mutation at np 3,196 was present. The normal sequence remained uncut at 617 np, while the mutant 16S^{3,196} was divided into 530 and 87 np fragments.

10

15

The ND2⁵⁴⁶⁰ G to A and G to T mutations were evaluated by several methods. To test for the G to A mutation, a 2820 np segment of mtDNA was amplified at $T_m=53^\circ\text{C}$ with primers at nps 3108-3127 \rightarrow and 5898-5917 plus a 10 np tail \leftarrow or a 332 np segment of mtDNA at $T_m=49^\circ\text{C}$ with primers at nps 5317-5333 \rightarrow and 5629-5648 \leftarrow . Digestion of the 2820 np segment with HphI gave 13 fragments ranging in size from 12 to 1130 np. A 491 np fragment located at nps 5315-5806 contained np 5460 and remains uncut with the Cambridge sequence, but was cut into 346 and 145 np fragments when the G to A nucleotide change was present. When the 332 np mtDNA segment was digested with HphI, it remained uncut when the Cambridge sequence was present and was cut into 198 and 134 np fragments when the G to A mutation was present (Figure 13). The second method was specific for the G to T mutation at np 5460. A primer was constructed with a mismatched "G" at np 5458 that created an AluI site only when the mutant T at np 5460 was present (mismatch primer sequence at nps 5441-5459: 5'-ATTCCTCCCCCACTCA*G*C-3' [SEQ ID No:1]). A 487 np segment of mtDNA was amplified at $T_m=49^\circ\text{C}$ with the mismatch primer \rightarrow and a primer at nps 5898-5917 plus a 10 np tail \leftarrow . Digestion with AluI produced fragments with sizes of 144, 104, 82, 60, 53, and 44 np for the Cambridge sequence and sizes of 125, 104, 82, 60, 53, 44, and 19 if the T at np 5460 was present. The third method

20

25

30

35

was specific for the Cambridge sequence G at np 5460. A primer was constructed with a mismatched "C" at np 5458 that created an *Aci*I site only when the Cambridge sequence "G" at np 5460 was present (mismatch primer sequence at nps 5441-5459: 5'-ATTCTCTCCCACACTCA"C"C-3' [SEQ ID NO:2]). A 208 np segment of mtDNA was amplified at $T_H=49^\circ\text{C}$ with the mismatch primer \rightarrow and a primer at nps 5629-5648 \leftarrow (Figure 13) or a 487 np segment of mtDNA was amplified at $T_H=49^\circ\text{C}$ with the mismatch primer \rightarrow and a primer at nps 5898-5917 plus a 10 np tail \leftarrow . Digestion with *Aci*I of PCR segments containing the Cambridge sequence G at np 5460 produced fragments with sizes of 190 and 18 np for the 208 np segment and sizes of 278, 173, 18, 15, and 3 np for the 487 np segment. Any nucleotide change at np 5460 resulted in no cut in the 208 np fragment and fragments with sizes 296, 173, 15, and 3 for the 487 np segment.

Results

Phylogenetic analysis of mtDNA mutations in AD, AD+PD, and PD

To identify mtDNA mutations that may be associated with Alzheimer's disease and Parkinson's disease, we determined the restriction endonuclease site patterns (haplotypes) of the mtDNAs from a group of 71 Caucasian patients, consisting of 33 with AD, 30 with AD+PD, and 8 with PD, as well as 74 Caucasian controls. Each mtDNA genome was amplified in nine overlapping polymerase chain reaction (PCR) fragments and each fragment was digested with 14 restriction endonucleases. Restriction site changes were detected by agarose gel electrophoresis and the phylogenetic relationships of the patient and control mtDNA haplotypes determined by the parsimony program PAUP^{18,42,53,54}. Novel restriction site variants were identified by comparing the patients' restriction site variants with those of our global data base for mtDNA restriction site polymorphisms⁵⁵.

Nine patients (two with AD, five with AD+PD, and two with PD) and one control were found to harbor an *Rsa* I site loss in the mtDNA D-loop at np 16,304, a region with no known function, and an *Ava* II site gain at np 4,336 which occurs in the tRNA^{Gln} gene (tRNA^{Gln}_{4,336}). The combination of these mutations caused these patients to cluster on

one discrete branch of the Caucasian mtDNA phylogeny (Figure 1; bold lines). Within this mtDNA^{16,304+4,336} lineage, two additional novel mtDNA mutations were found. One of these was an Rsa I site gain in ND1 at np 3,397 found in patient ADPD1. This same mutation was also found in patient ADPD2 outside the mtDNA^{16,304+4,336} lineage (Figure 1, enclosed by a box). The second novel mutation found in the mtDNA^{16,304+4,336} lineage was an insertion of approximately five nucleotides between nps 956 and 965 in the 12S rRNA gene of patient ADPD29. A third novel mutation was found outside the mtDNA^{16,304+4,336} lineage in patient AD10 (Figure 1, enclosed by a box). This individual harbored a heteroplasmic Dde I site loss at np 3,196 in the 16S rRNA gene.

The tRNA^{Gln4,336} mutation

To substantiate the clustering of the nine AD and PD mtDNAs within the mtDNA^{16,304+4,336} lineage, we sequenced 344 nps of the D-loop hypervariable region from each patient. Consistent with their forming a single mtDNA lineage, all were found to harbor the T to C transition at np 16,304, thus eliminating the Rsa I site. Two mtDNAs, ADPD28 and AD23, also had a second T to C transition at np 16,311 which eliminated an adjacent Rsa I site and ADPD28 also had a T to C transition mutation at np 16,172. Thus, all nine patients had very similar D-loop sequences, confirming that they belonged to the same mtDNA lineage. Divergence calculations using the observed D-loop and restriction site polymorphisms suggest that the tRNA^{Gln4,336} mutation arose in a Caucasian mtDNA 8,500 to 17,000 years ago.

To investigate the relationship of the tRNA^{Gln4,336} mutation to Alzheimer's disease and Parkinson's disease, we sequenced the tRNA^{Gln} gene from three of the patient mtDNAs. This revealed that all harbored the A to G transition mutation at np 4,336 necessary to create the Ava II site (Figures 2 and 3). This nucleotide connects the amino acid acceptor stem with the TΨC stem and is conserved in human, cow, and chicken mtDNAs, but not in mouse and *Xenopus* (Figure 4). The mutation was present in 3.2% (2/62) of brains with Alzheimer's disease, 6.8% (5/73) of brains with AD+PD, and 5.3% (2/38) of patients who were clinically defined as having Parkinson's disease,

giving an overall total of 5.2% (9/173) of the patients examined in this study (Table I). By contrast, the tRNA^{Gln4,336} mutation was present in 0.2% (1/547) of individuals analyzed in our population studies (laboratory controls) (see Table I legend for references).

5

Additional control data was obtained from published population surveys of mtDNA restriction site variants in which Ava II digestion was included (literature controls) (see Table I legend for references). This analysis revealed that this Ava II site gain has not been observed in Africans, Asians, or American Indians, but was reported infrequently in Caucasians (Table I). Since virtually all of the AD, AD+PD, and PD patients studied were Caucasian and the tRNA^{Gln4,336} mutation arose in a Caucasian mtDNA lineage, we compared the patient frequency of this Ava II site gain with that of all Caucasians surveyed to date. This analysis yielded a control frequency of 0.7% (12/1691) which includes 125 laboratory controls and 1566 literature controls. Compared to this frequency, the tRNA^{Gln4,336} mutation was 4.6 times more common in AD patients, 9.7 times more common in AD+PD patients, 7.6 times more common in PD patients, and 7.4 times more common in all AD and PD patients than in the Caucasian controls. While the number of patients who harbored the tRNA^{Gln4,336} mutation for the Alzheimer's disease (2/62) and PD (2/38) groups was too small to insure valid statistical comparisons, the AD+PD patients (5/73) and the combination of all AD, AD+PD, and PD patients (9/173) showed a statistically significant difference from the frequency of the tRNA^{Gln4,336} mutation observed in the total Caucasian control group at the $p < 0.005$ level. Similarly, comparison of the AD+PD or the total patient groups with all controls examined (12/3138 or 0.4%) was significant at $p < 0.005$. Thus, individuals harboring the tRNA^{Gln4,336} mutation appear to be at an increased risk for developing AD, AD+PD, or PD.

20

25

30

35

The observed frequency difference between the Caucasian patients and controls was not due to a sampling bias in which a subpopulation with an increased frequency of the tRNA^{Gln4,336} mutation was over-represented in our patient population, since we observed this mutation at increased frequency in patients from three different regions of the

United States: Massachusetts 2/68 (2.9%), Georgia and North Carolina 4/82 (4.8%), and southern California 3/23 (13%). Moreover, the Caucasian controls analyzed in our laboratory and those from the literature included random Californians and Georgians, British, Finnish, Italians, Jews, Arabs and Hindus, and none of these populations had significantly increased frequencies of tRNA^{Gln4,336} mutation relative to the population average (Table I, legend). Consequently, if a sampling bias did exist for our patients, it occurred independently in each of three different regions of the United States, and the increased input of the tRNA^{Gln4,336} mtDNAs would have had to come from an immigrant population which was a minor contributor to American colonization. Therefore, it seems most likely that the increased frequency of the tRNA^{Gln4,336} mutation in our Alzheimer's disease and Parkinson's disease patients was due to increased pathology rather than sampling error.

The ND1^{3,397[Met→Val]} mutation

To characterize the Rsa I site gain in ND1 which was found in ADPD1 within the mtDNA^{16,304+4,336} lineage and ADPD2 located outside this lineage, we sequenced this region of the mtDNA from these patients as well as from three controls. An A to G transition mutation was found at np 3,397 in all cases (Figures 5 and 6). This mutation substituted a valine for a methionine at the thirty-first amino acid in the ND1 protein. This methionine has been conserved throughout animal Oligomera evolution, extending down to fungi and some protozoa. It diverges in the invertebrate Polymera lineage and in plants (Figure 7, shaded area). Further, the ND1^{3,397[Met→Val]} mutation alters a highly conserved methionine within a hydrophilic block of 36 amino acids that is conserved from Trypanosoma brucei to man⁵⁶. Thus, this mutation could be functionally significant.

This mutation was found in two of the 73 AD+PD brains examined (2.7%). It was not found in 62 AD brains, 38 blood samples from clinically ascertained Parkinson's disease patients, or 11 brain samples from patients with other CNS diseases (Table II). Since the two AD+PD patients which harbor the ND1^{3,397[Met→Val]} were Caucasian, we

investigated the frequency of this mutation in Caucasian controls. No positives were found in 90 Caucasian laboratory controls or in 158 Caucasian literature controls (0/248 total Caucasian controls) (Table II). Thus, in our Caucasian sample this mutation was confined to the two AD+PD patients.

To clarify the genetic relationship between the two ND1^{3,397[Met→Val]} AD+PD patients, their mtDNA haplotypes were analyzed. The ADPD1 and ADPD2 mtDNAs differed by seven previously described restriction site variants⁵⁵, which placed them on two separate branches of the Caucasian mtDNA phylogeny (Figure 1). To further substantiate the wide divergence of these two mtDNAs, the entire D-loops from ADPD1 and ADPD2 were sequenced and found to differ by five nucleotide substitutions (nps 16270, 16304, 73, 150, 152) and by a dinucleotide deletion between nps 514 and 523 (Figure 8). Moreover, the D-loop of the "Cambridge sequence"⁵⁷ fell midway between the two patient sequences. Since the Cambridge sequence is known to have an A at np 3,397 while both patients are G, this analysis demonstrates that the ND1^{3,397[Met→Val]} mutation must have occurred two independent times (Figure 8). Since these independent events were associated with the same AD+PD neuropathology, it is possible that this mutation contributes to the pathogenesis of this disease.

Available pedigree information from these two individuals suggests that the ND1^{3,397[Met→Val]} mutation may be associated with the maternal transmission of AD. This mutation was present in the blood of the unaffected brother of ADPD2, confirming its maternal inheritance, and a history of cognitive decline was obtained for the mother of ADPD1 and for two maternal aunts of ADPD2.

Extending the population analysis to other ethnic groups showed that the mutation was not found in 63 Africans. However, it was found in 1/207 (0.5%) Asians and in 6/266 (2.3%) American Indians. All six of the American Indian positives were found in a sample of 28 Ticuna, a small, isolated tribe from the upper Amazon. In these six Ticuna, the mutation was found on only two almost identical mtDNA haplotypes which differed by one restriction site and were classified as

belonging to the American Indian specific mtDNA haplotype cluster D⁵⁴. Consequently, all six Ticuna represent a single founding mutational event that occurred within this isolated tribe. By contrast, the Asian mutation (an individual from Tibet) occurred on an Asian-specific haplotype⁵³ which is very different from those observed in the Ticunas (Table II legend). These results indicate that the Ticuna, the Asian, and the two Caucasian mutations are due to four independent mutational events.

In conclusion, the evolutionary conservation of the ND1^{3,397[Met→Val]} mutation, its presence in Caucasian patients but not Caucasian controls, and its association with affected maternal relatives suggests that this mutation could contribute to AD+PD pathology. However, the low frequency of the mutation in patients and its occasional appearance in the Asian and American Indian populations also raises the possibility that it is a neutral sequence variant.

The 12S^{956-965ins} mutation

One patient (ADPD29), within the mtDNA^{16,304+4,336} lineage, was found to harbor a novel 12S rRNA gene insertion (Figure 1). This insertion increased the size of the encompassing restriction fragment generated by multiple restriction endonucleases (e.g. Alu I, Dde I, Mbo I, etc.) (Figure 9). Direct sequence analysis revealed that the insertion consisted of approximately five cytosines within the sequence CCCCTCCCC [SEQ ID NO:3] between nps 956 to 965 (12S^{956-965ins}) (Figure 10). This insertion was not observed in 699 controls including 119 Caucasians^{42,58}, 241 Asians^{53,58}, 61 Africans⁵⁸, 167 American Indians⁵⁴, 90 New Guineans^{58,59}, or 21 Australian Aborigines⁵⁸, and is the first report of an insertion within a human mtDNA gene. The unique nature of this AD+PD mutation, its absence from the general population, and its association with both the tRNA^{6194,336} mutation and the AD+PD phenotype suggest that it may be functionally related to AD+PD.

The heteroplasmic 16S^{3,196} mutation

One patient (AD10) (Figure 1), who lies outside the mtDNA^{16,304+4,336} lineage, was found to harbor a heteroplasmic Dde I site loss. This mutation was found in approximately 50% of the mtDNAs of this individual's frontal, temporal, and parietal lobes as well as his basal ganglia and cerebellum. The mutation was sequenced and found to be the result of a G to A transition in the 16S rRNA gene at np 3,196 (16S^{3,196}) which also creates an Ssp I site (Figure 11).

To determine the frequency of this mutation in the general population, 699 individuals were surveyed (see the 16S^{956-965ins} section for references). Two individuals were found to have lost this Dde I site, an American Indian from the Pima tribe⁵⁴ and an Asian from Borneo⁵³. Direct sequencing revealed that the American Indian was homoplasmic for a C to T transition at np 3,192, while the Asian was homoplasmic for the same np 3,196 mutation as found in AD10 (Figure 12). Consistent with this result, the Ssp I site was present in the Asian, but absent in the American Indian (Figure 11). Since the 16S^{3,196} mutation was heteroplasmic in the Alzheimer's disease patient and homoplasmic in the control, the relevance of this mutation to the Alzheimer's disease neuropathology is unclear.

The ND2⁵⁴⁶⁰ mutations

Recently, it was reported that a high proportion of Alzheimer's disease patients harbored one of two mtDNA mutations in ND2 at np 5460: a G to A transition converting an alanine to threonine and a G to T transversion converting an alanine to serine⁶⁰. To determine the relevance of these mutations to our patients, we employed two restriction endonuclease tests (Figure 13). The G to A mutation was detected by an Hph I site gain, in which the 332 np PCR fragment was cut into 198 and 134 bp fragments. To test for either the G to A or the G to T mutations at np 5460, we prepared a PCR primer in which the second nucleotide from the 3'-end (np 5458) was changed from a T to C. This created an Aci I site in normal individuals having a G at np 5460, but eliminated the Aci I site when any other nucleotide was

substituted at np 5460. In normal individuals, the resulting 208 np PCR product was cut into 190 and 18 np fragments, but in mutants it remained uncut.

- 5 The G to A mutation was found in 4.4% (3/68) of patients with AD, AD+PD, and PD. However, it was also present in 8.6% (3/35) of Caucasian controls. Consequently, this mutation does not appear to be specific for Alzheimer's disease or Parkinson's disease and probably is a relatively common polymorphism. The G to T mutation was not
10 detected in any of 67 AD, AD+PD, or PD patients or in 41 controls. Hence, mutations at np 5460 do not appear to play a major role in our AD, AD+PD, or PD patients.

TABLE I

AD Patients	AD + PD Patients	PD Patients	Total Patients	Caucasian Laboratory Controls	Caucasian Literature Controls	Total Caucasian Controls	Total Laboratory Controls	Total Literature Controls	Total Controls
3.2%	6.8%	5.3%	5.2%	0.8%	0.7%	0.7%	0.2%	0.4%	0.4%
2/62	5/73	2/38	9/173	1/125	11/1566	12/1691	1/547	12/2591	12/3036

Table I. Association between the tRNA^{Gln} mutation and AD, AD+PD, and PD. Patients: The AD and AD+PD samples included 62 and 73 Caucasians, respectively. The PD group had 35 Caucasians, 1 Afro-American, and 2 Asians. Controls: The Laboratory Controls included the following groups. Caucasians of mixed European origin: 125; Africans: 44 (41 Senegalese and 3 Afro-Americans); American Indians: 167⁵⁴; Unknown ethnic origin: 4. The Literature Controls included the following groups. Caucasians: a group of 50 Europeans⁵⁵, a group of 46 Europeans⁵⁶, a group of 46 Italian Albanians (one with the Avail site gain)¹⁰¹, 75 Israeli Arabs and Jews (one with the Avail site gain)¹⁰², 100 Italian Apulians (one with the Avail site gain)¹⁰³, 87 Italian Apulians¹⁰⁴, 90 Italian Sicilians¹⁰⁵, 231 Italians (two with the Avail site gain)¹⁰⁶, 480 Italians (two with the Avail site gain)¹⁰⁷, 79 Indian Hindus¹⁰⁸, 110 Finnish (two with the Avail site gain)¹⁰⁹, 73 British (one Avail site gain)^{110,111}, 39 Jews¹¹², and 60 Italian Calabrians¹¹³; Africans and Afro-Americans: 74 Bantus and Bushmen⁵⁷, 20 Africans and Afro-Americans⁵⁸, and 186 Senegalese¹¹⁴; Asians: 46 Chinese and Japanese⁵⁹, 34 Asians⁶⁰, 122 Japanese¹¹⁵, 116 Japanese¹¹⁶, 121 Asians¹¹⁷ and 91 Tharu¹¹⁸; American Indians: 30 Warao⁶¹ and 74 Pima and Papago¹¹⁹; Australian aborigine and New Guineans: 21 aboriginals⁶², 26 Papua New Guineans⁶³ and 64 Papua New Guineans⁶⁴ who were mixed highland and lowland inhabitants. The Total Caucasian Controls included the Literature Controls plus the 125 individuals tested for this study (Laboratory Controls).

TABLE II

AD Patients	AD + PD Patients	PD Patients	Total Patients	Caucasian Laboratory Controls	Caucasian Literature Controls	Total Caucasian Controls	Total Laboratory Controls	Total Literature Controls	Total Controls
0%	2.7%	0%	1.2%	0%	0%	0%	*0.3-1.1%	0%	0.7%
0/62	2/73	0/38	2/173	0/90	0/158	0/248	2-7/612	0/439	7/1051

Table II. Association between the ND1³³⁹⁷ mutation and AD+PD. Patients: The AD and AD+PD samples included 62 and 73 Caucasians, respectively. The PD group had 35 Caucasians, 1 Afro-American, and 2 Asians. Controls: The Laboratory Controls included the following groups. Caucasians: 64 individuals analyzed for this study and 26 previously reported individuals². Asians: 153 Asians⁵³ and 54 Tibetans. American Indians: 266^{54,120}. Africans: 41 Senegalese, 2 Afro-Americans. Unknown ethnic origin: 6. The Literature Controls included the following groups. Caucasians: 46 Europeans⁵⁸, 39 Jews¹¹², 73 British^{110,111}. Asians: 34 Asians⁵⁸, 116 Japanese¹¹⁶. Africans: 20⁵⁸. Australian Aborigines: 21⁵⁸. Papua New Guineans: 26⁵⁸ and 64⁵⁹. The prevalence estimates for the ND1³³⁹⁷ mutation are expressed as a range of values (0.3% - 1.1%). The 0.3% estimate represents the number of mutational events, thus lumping the 6 Ticuna Indians together. The 1.1% estimate includes the total number of positive controls, counting all six of the Ticuna as independent events. The two Ticuna and one Tibetan haplotype differ from the Cambridge sequence by the following polymorphisms: Ticuna haplotypes: (A) 3,397+ [RsaI], 4,769- [AluI], 5,176- [HincII], 7,025+ [AluI], 8,858+ [HhaI], 10,394+ [DdeI], 10,397+ [AluI], 13,702- [HaeIII], 14,199- [HincII], 14,268+ [HinfI]; (B) 3,397+ [RsaI], 4,769- [AluI], 5,176- [AluI], 7,025+ [AluI], 8,858+ [HhaI], 10,394+ [DdeI], 10,397+ [AluI], 13,702- [HaeIII], 14,199- [HincII], 14,268+ [HinfI], 14,368+ [HinfI], DdeI- [15,073]. Asian haplotype: 663+ [HaeIII], 3,397+ [RsaI], 4,769- [AluI], 7,025+ [AluI], 8,858+ [HhaI], 13,702- [HaeIII], 14,199- [HincII], 14,268+ [HinfI], 14,368+ [HinfI]. (+) = site gain and (-) = site loss relative to the published sequence 57.

TABLE III

		Patients	Controls
ND2 ⁵⁴⁶⁰	G→A	3/68 (4.4%)	3/35 (8.6%)
ND2 ⁵⁴⁶⁰	G→T	0/67 (0%)	0/41 (0%)

Table III: Distribution of the ND2⁵⁴⁶⁰ polymorphism in individuals with AD, AD+PD, and AD and controls.

References

1. Whitehouse, P.J. et al. *Science* 215, 1237-1239 (1982).
2. Chui, H.C., Teng, E.L., Henderson, V.W. & Moy, A.C. *Neurology* 35, 1544-1550 (1985).
3. Mayeux, R., Stern, Y. & Spanton, S. *Neurology*. 35, 453-461 (1985).
4. Ditter, S.M. & Mirra, S.S. *Neurology*. 37, 754-760 (1987).
5. Mirra, S.S. et al. *Neurology* 41, 479-486 (1991).
6. Hansen, L.A. et al. *Neurology* 40, 1-8 (1990).
7. Byrne, E.J., Lennox, G., Lowe, J. & Godwin-Austin, R. J. *Neurol. Neurosurg. Psychiatry*. 52, 709-717 (1989).
8. Perry, R.H., Irving, D., Blessed, G., Fairbairn, A. & Perry, E.K. *J. Neurol. Sci.* 95, 119-139 (1990).
9. Dickson, D.W. et al. *Acta Neuropathol.* 78, 572-84 (1989).
10. Crain, B.J. & Mirra, S.S. *Current Opinion Neurol. Neurosurg.* 921-925 (1990).
11. Wallace, D.C. *Science* 256, 628-632 (1992).
12. Wallace, D.C. *Ann. Rev. Biochem.* 61, 1175-1212 (1992).
13. Mizuno, Y. et al. *Biochem. Biophys. Res. Commun.* 163, 1450-1455 (1989).
14. Schapira, A.H.V. et al. *Lancet* 1, 1269 (1989).
15. Schapira, A.H.V. et al. *J. Neurochem.* 54, 823-827 (1990).
16. Parker, W.D., Boyson, S.J. & Parks, J.K. *Ann. Neurol.* 26, 719-723 (1989).
17. Bindoff, L.A., Birch-Machin, M.A., Cartledge, N.E.F., Parker, Jr., W.D. & Turnbull, D.M. *J. Neurol. Sci.* 104, 203-208 (1991).
18. Shoffner, J.M., Watts, R.L., Juncos, J.L., Torroni, A. & Wallace, D.C. *Ann. Neurol.* 30, 332-339 (1991).
19. Parker, W.D., Filley, C.M. & Parks, J.K. *Neurology* 40, 1302-1303 (1990).
20. Sims, N.R., Finegan, J.M., Blass, J.P., Bowen, D.M. & Neary, D. *Brain Res.* 436, 30-38 (1987).
21. Peterson, C & Goldman, J.E. *Proc. Natl. Acad. Sci. USA* 83, 2758-2762 (1986).

22. Blass, J.P., Baker, A.C., Ko, L. & Black, R.S. *Arch. Neurol.* 47, 864-869 (1990).
23. Shoffner, J.M. & Wallace, D.C. *Adv. Hum. Genet.* 19, 267-330 (1990).
24. Hansen, L.A., Masliah, E., Terry, R.D. & Mirra, S.S. *Acta Neuropathol.* 78, 194-201 (1989).
25. Saraiva, A.A., Borges, M.M., Madeira, M.D., Tavares, M.A. & Paula-Barbosa, M.M. *J. Submicrosc. Cytol.* 17, 459-64 (1985).
26. Giles, R.E., Blanc, H., Cann, H.M. & Wallace, D.C. *Proc. Natl. Acad. Sci. USA* 77, 6715-6719 (1980).
27. Case, J.T. & Wallace, D.C. *Somat. Cell Genet.* 7, 103-108 (1981).
28. Wallace, D.C. *Somat. Cell Mol. Genet.* 12, 41-49 (1986).
29. Lott, M.T., Voljavec, A.S. & Wallace, D.C. *Am. J. Ophthalmol.* 109, 625-631 (1990).
30. Shoffner, J.M. et al. *Proc. Natl. Acad. Sci. USA* 86, 7952-7956 (1989).
31. Shoffner, J.M. et al. *Cell* 61, 931-937 (1990).
32. Wallace, D.C. et al. *Cell* 55, 601-610 (1988).
33. Holt, I.J., Harding, A.E., Petty, R.K.H. & Morgan-Hughes, J.A. *Am. J. Hum. Genet.* 46, 428-433 (1990).
34. Trounce, I., Byrne, E. & Marzuki, S. *Lancet* 1, 637-639 (1989).
35. Yen, T.-C., Chen, Y.-S., King, K.-L., Yeh, S.-H. & Wei, Y.-H. *Biochem. Biophys. Res. Commun.* 165, 994-1003 (1989).
36. Muller-Hocker, J. *Am. J. Pathol.* 134, 1167-1173 (1989).
37. Cortopassi, G.A. & Arnheim, N. *Nucleic Acids Res.* 18, 6927-6933 (1990).
38. Corral-Debrinski, M. et al. *J. A. M. A.* 266, 1812-1816 (1991).
39. Ikebe, S. et al. *Biochem. Biophys. Res. Commun.* 170, 1044-1048 (1990).
40. Ozawa, T. et al. *Biochem. Biophys. Res. Commun.* 170, 830-836 (1990).
41. Wallace, D.C. et al. *Science* 242, 1427-1430 (1988).
42. Brown, M.D. et al. *Genetics* 130, 163-173 (1992).
43. Newman, N.J. & Wallace, D.C. *Am. J. Ophthal.* 109, 727-730 (1990).

44. Brown, M.D., Voljavec, A.S., Lott, M.T., MacDonald, I. & Wallace, D.C. *FASEB Journal* 6,2791-2799 (1992).
45. Johns, D.R. & Berman, J. *Biochem. Biophys. Res. Commun.* 174, 1324-1330 (1991).
46. Johns, D.R. & Neufeld, M.J. *Biochem. Biophys. Res. Commun.* 181, 1358-1364 (1991).
47. Vilkki, J., Ott, J., Savontaus, M.L., Aula, P. & Nikoskelainen, E.K. *Am. J. Hum. Genet.* 48, 486-491 (1991).
48. Howell, N., Kubacka, I., Yu, M. & McCullough, D.A. *Am. J. Hum. Genet.* 48, 935-942 (1991).
49. Howell, N. et al. *Am. J. Hum. Genet.* 49, 939-950 (1991b).
50. Huoponen, K., Vilkki, J., Aula, P., Nikoskelainen, E.K. & Savontaus, M.L. *Am. J. Hum. Genet.* 48, 1147-1153 (1991).
51. Brown, M.D. et al. *Am. J. Hum. Genet.* 51,378-385 (1992).
52. Singh, G., Lott, M.T. & Wallace, D.C. *N. Engl. J. Med.* 320, 1300-1305 (1989).
53. Ballinger, S.W. et al. *Genetics* 130, 139-152 (1992).
54. Torroni, A. et al. *Genetics* 130, 153-162 (1992).
55. Wallace, D.C., Lott, M.T., Torroni, A. & Shoffner, J.M. *Cytogenet. Cell Genetic.* 58, 1103-1123 (1991).
56. Burger, G. & Werner, S. *J. Mol. Biol.* 186, 231-242 (1985).
57. Anderson, S. et al. *Nature* 290, 457-465 (1981).
58. Cann, R.L., Stoneking, M. & Wilson, A.C. *Nature* 325, 31-36 (1987).
59. Stoneking, M., Jordem, L.B., Bhatia, K. & Wilson, A.C. *Genetics* 124, 717-733 (1990).
60. Lin, F.-H. et al. *Biochem. Biophys. Res. Commun.* 182, 238-246 (1992).
61. Farrer, L.A., Myers, R.H., Connor, L., Cupples, L.A. & Growdon J.H. *Am. J. Hum. Genet.* 48, 1026-1033 (1991).
62. Haines, J. *Am. J. Hum. Genet.* 48, 1021-1025 (1991).
63. Pericak-Vance, M.A. et al. *Am. J. Hum. Genet.* 48, 1034-1050 (1991).
64. Chartier-Harlin, M.C. et al. *Nature* 353, 844-846 (1991).
65. Goate, A. et al. *Nature* 349, 704-706 (1991).

66. Murrell, J., Farlow, M., Ghetti, B. & Benson, M.D. *Science* 254, 97-99 (1991).
67. Naruse, S. et al. *Lancet* 337, 978-979 (1991).
68. Yoshioka, K., Miki, T., Katsuya, T., Ogihara, T. & Sakaki, Y. *Biochem. Biophys. Res. Commun.* 178, 1141-1146 (1991).
69. Tanzi, R.E. et al. *Nature* 329, 156-157 (1987).
70. Van Broeckhoven, C. et al. *Nature* 329, 153-155 (1987).
71. Johnson, W.G. *Neurology* 41, 82-87 (1991).
72. Golbe, L.I. *Neurology* 40, 7-14 (1990).
73. Goto, Y.-i., Tojo, M., Tohyama, J., Horai, S., & Nonaka, I. *Ann. Neurol.* 31, 672-675 (1992).
74. Goto, Y., Nonaka, I. & Horai, S. *Nature* 348, 651-653 (1990).
75. Kobayashi, Y. et al. *Biochem. Biophys. Res. Commun.* 173, 816-822 (1990).
76. Kosik, K.S. *Science* 256, 780-783 (1992).
78. Zeviani et al., *Lancet* 338, 143-147 (1991).
79. Singer, T.P. & Ramsey, R.R. *FEBS Lett.* 274, 1-8 (1990).
80. Beal, M.F. *Ann. Neurol.* 31, 119-130 (1992).
81. Khachaturian, Z.S. *Arch. Neurol.* 42, 1097-1105 (1985).
82. Wilkinson L. *SYSTAT: The System for Statistics*. Evanston, IL: SYSTAT, Inc 1989:142-179, 462-463.
83. Johnson, M.J., Wallace, D.C., Ferris, S.D., Rattazi, M.C. & Cavalli-Sforza, L.L. *J. Mol. Evol.* 19, 255-271 (1983).
84. Cann, R.L., Brown, W.M. & Wilson, A.C. *Genetics* 106, 479-499 (1984).
85. Swofford, D.L. *PAUP: Phylogenetic analysis using parsimony*, version 3.0. Computer program distributed by the Illinois Natural History Survey, Champaign, Ill (1990).
86. Nei, M. & Tajima, F. *Genetics* 105, 207-217 (1983).
87. Vigilant, L., Pennington, R., Harpending, H., Kocher, T.D. & Wilson, A.C. *Proc. Natl. Acad. Sci. USA* 86, 9350-9354 (1989).
88. Anderson, S. et al. *J. Mol. Biol.* 156, 683-717 (1982).
89. Bibb, M.J., Van Etten, R.A., Wright, C.T., Walberg, M.W. & Clayton, D.A. *Cell* 26, 167-180 (1981).

90. Gadaleta, G. et al. *J. Mol. Evol.* 28, 497-516 (1989).
91. Desjardins, P. & Morais, R. *J. Mol. Biol.* 212, 599-634 (1990).
92. Roe, B.A., Ma, D.-P., Wilson, R.K. & Wong, J.F.-H. *J. Biol. Chem.* 17, 9759-9774 (1985).
93. Jacobs, H.T., Elliott, D.J., Veerabhadracharya, B.M. & Farquharson, A. *J. Mol. Biol.* 202, 185-217 (1988).
94. Clary, D.O. & Wolstenholme, D.R. *J. Mol. Evol.* 22, 252-271 (1985).
95. Okimoto, R., Macfarlane, J.L., Clary, D.O. & Wolstenholme, D.R. *Genetics* 130, 471-498 (1992).
96. Pritchard, A.E., Sable, C.L., Venuti, S.E. & Cummings, D.J. *Nucleic Acids Res.* 18, 163-171 (1990).
97. Cummings, D.J., Domenico, J.M., & Michel, F. *Curr. Genet.* 14, 253-264 (1988).
98. Boer, P.H. & Gray, M.W. *EMBO J.* 7, 3501-3508 (1988).
99. Zwieb, C., Glotz, C. & Brimacombe, R. *Nucleic Acids Res.* 9, 3621-3640 (1981).
101. Dionne et al., *Med. Sci. Sports Exerc.* 23, 177-185 (1991).
100. Glotz, C., Zwieb, C. & Brimacombe, R. *Nucleic Acids Res.* 9, 3287-3306 (1981).
102. Bonne-Tamir, B., Johnson, M.J., Natali, A., Wallace, D.C. & Cavalli Sforza, L.L. *Am. J. Hum. Genet.* 38, 341-351 (1986).
103. Torroni, A. et al. *Int. J. Anthropol.* 5, 97-104 (1990).
104. De Benedictis, G., Rose, G., Passarino, G. & Quagliariello, C. *Ann. Hum. Genet.* 53, 311-318 (1989).
105. Semino, O. et al. *Ann. Hum. Genet.* 53, 193-202 (1989).
106. Brega, A. et al. *Ann. Hum. Genet.* 50, 327-338 (1986).
107. Sartoris, S. et al. *Ann. Hum. Genet.* 52, 327-340 (1988).
108. Semino, O., Torroni, A., Scozzari, R., Brega, A. & Santachiara Benerecetti, A.S. *Ann. Hum. Genet.* 55, 123-136 (1991).
109. Vilkkki, J., Savontaus, M.L. & Nikoskelainen, E.K. *Hum. Genet.* 80, 317-321 (1988).
110. Holt, I.J., Harding, A.E. & Morgan-Hughes, J.A. *Hum. Genet.* 79, 53-57 (1988).

111. Holt, I.J., Miller, D.H. & Harding, A.E. *J. Neurol. Neurosurg. Psychiatry* 51,1075-1077 (1988).
112. Tikochinski, Y., Ritte, U., Gross, S.R., Prager, E.M. & Wilson, A.C. *Am. J. Hum. Genet.* 48, 129-136 (1991).
113. De Benedictis, G., Rose, G., Caccio, S., Picardi, P., & Quagliariello, C. *Gene. Geogr.* 3, 33-40 (1989).
114. Scozzari, R. et al. *Am. J. Hum. Genet.* 43, 534-544 (1988).
115. Harihara, S., Hirai, M. & Omoto, K. *Japanese J. Hum. Genet. [Jinrui Idengaku Zasshi]* 31, 73-83 (1986).
116. Horai, S. & Matsunaga, E. *Hum. Genet.* 72, 105-117 (1986).
117. Harihara, S. et al. *Am. J. Hum. Genet.* 43, 134-143 (1988).
118. Brega, A. et al. *Am. J. Hum. Genet.* 39, 502-512 (1986).
119. Wallace, D.C., Garrison, K. & Knowler, W.C. *Am. J. Phys. Anthropol.* 68, 149-155 (1985).
120. Schurr, T.G. et al. *Am. J. Hum. Genet.* 46, 613-623 (1990).
121. Majander, A., Huoponen, K., Savontaus, M.-L., Nikoskelainen, E., & Wikstrom, M. *FEBS Letters* 292,289-292 (1991).
122. Larsson, N.-G. et al. *Am. J. Hum. Genet.* 50,360-363 (1992).
123. Parker, W.D., Oley, C.A., & Parks, J.K. *Lancet* 320,1331-1333 (1989).
124. Maniatis et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
125. Berger and Kimmel (1987) "Guide to Molecular Cloning Techniques," *Methods in Enzymology*, Volume 152, Academic Press, Inc., San Diego, CA.
126. Gibbs et al. (1989) *Nucleic Acids Res.* 17:2437.
127. Kwok et al. (1990) *Nucleic Acids Res.* 18:999.
128. Miyada et al. (1987) *Methods Enzymol.* 154:94.
129. Anderson et al., *Nature* 290:457-465 (1981).
130. DiHella et al. (1988) *Lancet* 1:497.
131. *Promega Protocols and Applications Guide* (1991) Promega Corporation, Madison, Wisconsin.
132. Orita et al. (1989) *Genomics* 5:874-879.
133. Orita et al. (1990) *Genomics* 6:271-276.

134. Newton et al. (1989) *Nucleic Acids Res.* 17:2503.
135. Nichols et al. (1989) *Genomics* 5:535.
136. Okayama et al. (1989) *J. Lab. Clin. Med.* 114:105.
137. Sarkar et al. (1990) *Anal. Biochem.* 186:64.
138. Sommer et al. (1989) *Mayo Clin. Proc.* 64:1361.
139. Wu (1989) *Proc. Natl. Acad. Sci. (U.S.A.)* 86:2757
140. Dutton et al. (1991) *Biotechniques* 11:700.
150. Baany et al. (1991) *Proc. Natl. Acad. Sci. (U.S.A.)* 88:189
151. R.A. Weiss (1991) *Science* 254:1992.
152. Frohman and Martin, *Cell* (1989) 56:145.
153. Wirak et al., the *EMBO Journal*, 10(2) 289-296 (1991).
154. Schilling et al., *Gene* 98(2) 225-230 (1991).
155. Quon, et al. (1991) *Nature* 352:239.
156. Wirak, et al. (1991) *Science* 253:323.
157. Kawabata, et al. (1991) *Nature* 354:476.
158. Mendelson et al. *Virology* 166:154-165.
159. Wondisford et al. (1988) *Molec. Endocrinol.* 2:32-39 (1988).
160. Kohler and Milstein, (1975) *Nature* 256:495-497.
161. *Antibodies: A Laboratory Manual*, Harlow and Lane (1988), Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
162. Winacker (1987) "From Genes to Clones," VCH Publishers, New York, NY).
163. Quenn et al. (1986) *Immunol. Rev.* 89:49-68.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Emory University School of Medicine
- (B) STREET: 1440 Clifton Road, N.E.
- (C) CITY: Atlanta
- (D) STATE: Georgia
- (E) COUNTRY: United States of America
- (F) POSTAL CODE (ZIP): 30322
- (G) TELEPHONE: 404/727-2211
- (H) TELEFAX: 404/727-2485
- (I) TELEX: none

(ii) TITLE OF INVENTION: "DETECTION OF MITOCHONDRIAL DNA MUTATIONS ASSOCIATED WITH ALZHEIMER'S DISEASE AND PARKINSON'S DISEASE"

(iii) NUMBER OF SEQUENCES: 3

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 EPO

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: Not yet available

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 07/963,723
- (B) FILING DATE: 20-OCT-1992

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATTCTCCCC AACTCAGC

19

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid

50

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATTCCTCCCC AACTCACC

19

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCCCCTCCCC

10

What is claimed is:

1. A method of diagnosing or predicting a predisposition to Alzheimer's disease and/or Parkinson's disease, comprising detecting in a sample from a subject the presence of a mutation of mitochondrial DNA which correlates to Alzheimer's disease and/or Parkinson's disease, the presence of the mutation indicating the presence of or a predisposition to Alzheimer's disease and/or Parkinson's disease.
2. The method of claim 1, wherein the mutation is in nucleotide position 4,336 of mitochondrial DNA.
3. The method of claim 2, wherein the detecting step comprises combining a nucleotide probe capable of selectively hybridizing to a nucleic acid containing the mutation with a nucleic acid in the sample and detecting the presence of hybridization.
4. The method of claim 2, wherein the detecting step comprises amplifying the nucleotides of the mutation and detecting the presence of the mutation in the amplified product.
5. The method of claim 2, wherein the detecting step comprises selectively amplifying the nucleotides of the mutation and detecting the presence of amplification.
6. The method of claim 2, wherein the detecting step comprises detecting the gain or loss of a restriction endonuclease site as detected by enzyme digest of the nucleotides of the mutation.
7. The method of claim 2, wherein the detecting step comprises utilizing Ligase Chain Reaction.
8. The method of claim 1, wherein the mutation is in nucleotide position 3,397 of mitochondrial DNA.
9. The method of claim 8, wherein the detecting step comprises combining a nucleotide probe capable of selectively hybridizing to a

nucleic acid containing the mutation with a nucleic acid in the sample and detecting the presence of hybridization.

10. The method of claim 8, wherein the detecting step comprises amplifying the nucleotides of the mutation and detecting the presence of the mutation in the amplified product.

11. The method of claim 8, wherein the detecting step comprises selectively amplifying the nucleotides of the mutation and detecting the presence of amplification.

12. The method of claim 8, wherein the detecting step comprises detecting the gain or loss of a restriction endonuclease site as detected by enzyme digest of the nucleotides of the mutation.

13. The method of claim 8, wherein the detecting step comprises utilizing Ligase Chain Reaction.

14. The method of claim 1, wherein the mutation is in nucleotide position 3,196 of mitochondrial DNA.

15. The method of claim 14, wherein the detecting step comprises combining a nucleotide probe capable of selectively hybridizing to a nucleic acid containing the mutation with a nucleic acid in the sample and detecting the presence of hybridization.

16. The method of claim 14, wherein the detecting step comprises amplifying the nucleotides of the mutation and detecting the presence of the mutation in the amplified product.

17. The method of claim 14, wherein the detecting step comprises selectively amplifying the nucleotides of the mutation and detecting the presence of amplification.

18. The method of claim 14, wherein the detecting step comprises detecting the loss of a restriction endonuclease site as detected by enzyme digest of the nucleotides of the mutation.

19. The method of claim 14, wherein the detecting step comprises utilizing Ligase Chain Reaction.
20. The method of claim 1, wherein the mutation is one or more inserted G-C base pairs between positions 956 and 965 of mitochondrial DNA.
21. The method of claim 20, wherein the detecting step comprises combining a nucleotide probe capable of selectively hybridizing to a nucleic acid containing the mutation with a nucleic acid in the sample and detecting the presence of hybridization.
22. The method of claim 20, wherein the detecting step comprises amplifying the nucleotides of the mutation and detecting the presence of the mutation in the amplified product.
23. The method of claim 20, wherein the detecting step comprises selectively amplifying the nucleotides of the mutation and detecting the presence of amplification.
24. The method of claim 20, wherein the detecting step comprises detecting the increase in size of a restriction fragment created by an enzyme digest of the nucleotides of the insertion.
25. The method of claim 20, wherein the detecting step comprises utilizing Ligase Chain Reaction.
26. A method of determining mutations associated with the presence of or predisposition to Alzheimer's disease and/or Parkinson's disease comprising:
- (a) obtaining a mitochondrial DNA-containing sample from a subject with Alzheimer's disease and/or Parkinson's disease;
 - (b) determining the presence of mutations in the mitochondrial DNA;
 - (c) comparing the mutations to mutations found in a normal subject; and

(d) determining which mutations have a greater rate of occurrence in the subject with Alzheimer's disease and/or Parkinson's disease.

27. A method of screening for an agent capable of treating Alzheimer's disease and/or Parkinson's disease comprising administering the agent to an animal having a mutation in mitochondrial DNA associated with Alzheimer's disease and/or Parkinson's disease and monitoring the effect of the agent on the animal's condition.

28. The method of claim 27, wherein the mutation in mitochondrial DNA is in nucleotide position 4,336.

29. The method of claim 27, wherein the mutation in mitochondrial DNA is in nucleotide position 3,397.

30. The method of claim 27, wherein the mutation in mitochondrial DNA is in nucleotide position 3,196.

31. The method of claim 27, wherein the mutation in mitochondrial DNA is one or more inserted G-C base pairs between positions 956 and 965 of mitochondrial DNA.

32. An isolated nucleic acid characteristic of human mitochondrial DNA, wherein the nucleic acid has a mutation at the nucleotide corresponding to position 4,336 of the mitochondrial DNA.

33. A structural RNA encoded by the nucleic acid of claim 32.

34. An isolated nucleic acid primer capable of selectively amplifying the mutation of claim 32.

35. An isolated nucleic acid of human mitochondrial DNA, wherein the nucleic acid has a mutation at the nucleotide corresponding to position 3,397.

36. A polypeptide encoded by the nucleic acid of claim 35.
37. An isolated nucleic acid primer capable of selectively amplifying the mutation of claim 35.
38. A method of diagnosing or predicting a predisposition to Alzheimer's Disease, comprising detecting in a sample from a subject the presence of a mutation of mitochondrial DNA which correlates to Alzheimer's Disease, the presence of the mutation indicating the presence of or a predisposition to Alzheimer's Disease.




FIG. 1A

SUBSTITUTE SHEET (RULE 26)

2/15




FIG. 1B




FIG. 2A

FIG. 2B

4/15




FIG. 3




FIG. 4




FIG. 5A

FIG. 5B

7/15




FIG. 6

8/15




FIG. 7

9/15




FIG. 8

10/15




FIG. 9A

FIG. 9B

FIG. 9C

11/15




FIG. 10

12/15

Dde I Site Loss




FIG. 11A

Ssp I Site Gain




FIG. 11B

13/15




FIG. 11C

14/15




FIG. 12




FIG. 13B

FIG. 13A

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/10072

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12Q 1/68; C12P 19/34; C07H 21/02, 21/04; C07K 13/00

US CL : 435/6, 91.2; 514/2, 44, 536/23.5, 24.31, 24.33; 530/350

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.2; 514/2, 44, 536/23.5, 24.31, 24.33; 530/350

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, BIOSIS

search terms: mitochondria, mutation, Alzheimer's, Parkinson's

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Bioch. Biophys. Res. Comm., Vol. 170, Number 3, issued 16 August 1990, Ikebe et al., "Increase of deleted mitochondrial DNA in the striatum in Parkinson's disease and senescence", pages 1044-1048.	1-38
A	JAMA, Vol. 266, Number 13, issued 02 October 1991, Randall, "Mitochondrial DNA: A new frontier in acquired and inborn gene defects", pages 1739-1740.	1-38

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T	later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A document defining the general state of the art which is not considered to be part of particular relevance	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E earlier document published on or after the international filing date	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* A	document member of the same patent family
* O document referring to an oral disclosure, use, exhibition or other means		
* P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

30 November 1993

Date of mailing of the international search report

JAN 31 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

KENNETH R. HORLICK

Facsimile No. NOT APPLICABLE

Telephone No. (703) 305-5126

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/10072

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Bioch. Biophys. Res. Comm., Vol. 176, Number 2, issued 30 April 1991, Ozawa et al., "Distinct clustering of point mutations in mitochondrial DNA among patients with mitochondrial encephalomyopathies and with Parkinson's Disease", pages 938-946.	1-38
A	Bioch. Biophys. Res. Comm., Vol. 186, Number 1, issued 15 August 1992, Petruzzella et al., "Is a point mutation in the mitochondrial ND2 gene associated with Alzheimer's disease?", pages 491-497.	1-38
A	Science, Vol. 256, issued 01 May 1992, Wallace, "Mitochondrial genetics: A paradigm for aging and degenerative diseases?", pages 628-632.	1-38
A	Biochem. Biophys. Res. Comm., Vol. 182, Number 1, issued 15 January 1992, Lin et al., "Detection of point mutations in codon 331 of mitochondrial NADH dehydrogenase subunit 2 in Alzheimer's brains", pages 238-246.	1-38